

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number  
**WO 2006/021002 A2**

(51) International Patent Classification:

**C07D 241/20** (2006.01) **A61K 31/4965** (2006.01)  
**C07D 401/04** (2006.01) **A61K 31/497** (2006.01)  
**C07D 403/04** (2006.01) **A61P 35/00** (2006.01)  
**C07D 413/04** (2006.01)

(21) International Application Number:

PCT/US2005/029518

(22) International Filing Date: 18 August 2005 (18.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/602,968 19 August 2004 (19.08.2004) US

(71) Applicant (for all designated States except US): **ICOS CORPORATION** [US/US]; 22021 20th Avenue S.E., Bothell, WA 98021 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FAROUIZ, Francine, S.** [FR/US]; 4222 93rd Avenue S.E., Mercer Island, WA 98040 (US). **HOLCOMB, Ryan** [US/US]; 1214 2nd Avenue, Kirkland, WA 98033 (US). **THORSETT, Eugene** [US/US]; 410 2nd Ave S., #304, Kirkland, WA 98033 (US). **GAUDINO, John, Joseph** [US/US]; 4224 Prairie Fire Circle, Longmont, CO 80503 (US).

(74) Agent: **NAPOLI, James, J.**; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, Suite 6300, Sears Tower, Chicago, IL 60606-6357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS USEFUL FOR INHIBITING CHK1

(57) Abstract: Aryl- and heteroaryl-substituted urea compounds useful in the treatment of diseases and conditions related to DNA damage or lesions in DNA replication are disclosed. Methods of making the compounds, and their use as therapeutic agents, for example, in treating cancer and other diseases characterized by defects in DNA replication, chromosome segregation, or cell division also are disclosed.



**WO 2006/021002 A2**

BEST AVAILABLE COPY

**COMPOUNDS USEFUL FOR INHIBITING CHK1****FIELD OF THE INVENTION**

The present invention relates to compounds useful for inhibiting enzymes that maintain and repair the integrity of genetic material. More particularly, the present invention relates to a series of aryl- and heteroaryl-substituted urea compounds, methods of making the compounds, and their use as therapeutic agents, for example, in treating cancer and other diseases characterized by defects in deoxyribonucleic acid (DNA) replication, chromosome segregation, or cell division.

**BACKGROUND OF THE INVENTION**

A large variety of diseases, conditions, and disorders (hereinafter "indications") are characterized as involving aberrantly proliferating cells. As used herein, "aberrantly proliferating cells" (or "aberrant cell proliferation") means cell proliferation that deviates from the normal, proper, or expected course. For example, aberrant cell proliferation includes inappropriate proliferation of cells wherein DNA or other cellular components have become damaged or defective. Aberrant cell proliferation also includes indications caused by, mediated by, or resulting in inappropriately high levels of cell division, inappropriately low levels of cell death (e.g., apoptosis), or both. Such indications can be characterized, for example, by single or multiple local abnormal proliferations of cells, groups of cells or tissue(s), and include cancerous (benign or malignant) and noncancerous indications.

- 2 -

By definition, all cancers (benign and malignant) involve some form of aberrant cell proliferation. Nonlimiting examples include carcinomas and sarcomas. Others are discussed below. Some noncancerous indications also involve aberrant cell proliferation. Nonlimiting examples of noncancerous indications involving aberrant cell proliferation include rheumatoid arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus. Others are discussed below.

One approach to treating indications involving aberrantly proliferating cells involves the use of DNA damaging agents. These agents are designed to kill aberrantly proliferating cells by disrupting vital cellular processes such as DNA metabolism, DNA synthesis, DNA transcription, and microtubule spindle formation. They also can operate, for example, by introducing lesions into DNA that perturb chromosomal structural integrity. DNA damaging agents are designed and administered in ways that attempt to induce maximum damage and consequent cell death in aberrantly proliferating cells with a minimum damage to normal, healthy cells.

A large variety of DNA damaging agents have been developed to date. Others are also in development. DNA damaging agents include chemotherapeutics and radiation. Unfortunately, the effectiveness of DNA damaging agents in treating conditions involving aberrant cell proliferation have been less than desired, particularly in the treatment of cancer. The selectivity of such agents for aberrantly proliferating cells over healthy cells (sometimes referred to as the therapeutic index) often is marginal.

Moreover, all cells have sensing and repair mechanisms that can work at cross purposes to DNA damaging agents. Such sensing mechanisms, called cell cycle

- 3 -

checkpoints, help to maintain the order of the various cell replication stages and to ensure that each step is executed with high fidelity (Hartwell et al., *Science*, 246:629-634 (1989); Weinert et al., *Genes Dev.*, 8:652 (1994)). When cells detect DNA damage, including damage purposefully induced by DNA damaging agents, certain signaling pathways activate cell cycle checkpoints and the cell replication cycle temporarily ceases ("arrests"). This arrest allows time for aberrantly proliferating cells to repair their DNA, often to a degree sufficient to allow the affected cells to continue to survive and proliferate. This unwanted repair tends to undermine efforts to induce DNA damage sufficient to kill aberrantly proliferating cells.

For example, the chemotherapeutic agent called Gemzar™ (gemcitabine, or 2',2' difluoro-2'-deoxycytidine) damages DNA by incorporating itself into DNA during synthesis. Left unrepaired, damaged DNA generally is rendered incapable of sustaining life. In many targeted cells, however, cell cycle checkpoints detect the improperly made (or otherwise damaged) DNA. The activated cell cycle checkpoints trigger cell cycle arrest for a time sufficient to allow damaged DNA to be repaired. This is one way in which aberrantly proliferating cells are theorized to resist the cell-killing effect of DNA-damaging agents such as chemotherapeutics, radiation, and other therapies.

Other DNA-damaging agents cause tumor cells to arrest in S-phase. Tumor cells have been observed to resist certain chemotherapeutics simply by arresting in S phase while the chemotherapeutic agent is being administered. Then, as soon as the drug is removed, DNA damage is repaired, cell cycle arrest ceases, and the cells progress through the remainder of the cell cycle (Shi et

- 4 -

al., *Cancer Res.* 61:1065-1072, 2001). Other therapeutics cause cell cycle arrest at other checkpoints, including G1 and G2 (described more fully below). DNA damaging agents that activate cell cycle checkpoints generally are referred to herein as "checkpoint activators." DNA damaging agents that activate the checkpoint designated "Chk1" (pronounced "check-one") are referred to herein as "Chk1 activators." Inhibitors of such checkpoints, generally and specifically, are referred to herein as "checkpoint inhibitors" and "Chk1 inhibitors," respectively.

Inhibition of various DNA damage checkpoints therefore is expected to assist in preventing cells from repairing therapeutically induced DNA damage and to sensitize targeted cells to DNA damaging agents. Such sensitization is in turn expected to increase the therapeutic index of these therapies.

To more fully understand the present invention, the following is a more detailed discussion of cell cycle phases and the role of Chk1.

The cell cycle is structurally and functionally the same in its basic process and mode of regulation across all eukaryotic species. The mitotic (somatic) cell cycle consists of four phases: the G1 (gap) phase, the S (synthesis) phase, the G2 (gap) phase, and the M (mitosis) phase. The G1, S, and G2 phases are collectively referred to as interphase of the cell cycle. During the G1 phase, biosynthetic activities of the cell progress at a high rate. The S phase begins when DNA synthesis starts, and ends when the DNA content of the nucleus of the cell has been replicated and two identical sets of chromosomes are formed.

The cell then enters the G2 phase, which continues until mitosis starts. In mitosis, the chromosomes

- 5 -

pair and separate, two new nuclei form, and cytokinesis occurs in which the cell splits into two daughter cells each receiving one nucleus containing one of the two sets of chromosomes. Cytokinesis terminates the M phase and marks the beginning of interphase of the next cell cycle. The sequence in which cell cycle events proceed is tightly regulated, such that the initiation of one cell cycle event is dependent on the completion of the prior cell cycle event. This allows fidelity in the duplication and segregation of genetic material from one generation of somatic cells to the next.

It has been reported that cell cycle checkpoints comprise at least three distinct classes of polypeptides, which act sequentially in response to cell cycle signals or defects in chromosomal mechanisms (Carr, A.M., *Science*, 271:314-315 (1996)). The first class is a family of proteins that detect or sense DNA damage or abnormalities in the cell cycle. These sensors include Ataxia-telangiectasia Mutated protein (Atm) and Ataxia-telangiectasia Rad-related protein (Atr). The second class of polypeptides amplify and transmit the signal detected by the detector and is exemplified by Rad53 (Alen et al. *Genes Dev.* 8:2416-2488 (1994)) and Chk1. A third class of polypeptides includes cell cycle effectors, such as p53, which mediate a cellular response, for example, arrest of mitosis and apoptosis.

Much of the current understanding of the function of cell cycle checkpoints has been derived from the study of tumor derived cell lines. In many cases, tumor cells have lost key cell cycle check points (Hartwell et al., *Science* 266:1821-28, 1994). It has been reported that a key step in the evolution of cells to a neoplastic state is the acquisition of mutations that inactivate cell cycle checkpoint pathways, such as those involving

- 6 -

p53 (Weinberg, R.A., *Cell* 81:323-330, 1995; Levine, A. J., *Cell* 88:3234-331, 1997). Loss of these cell cycle checkpoints results in the replication of tumor cells despite DNA damage.

5 Noncancerous tissue, which has intact cell cycle checkpoints, typically is insulated from temporary disruption of a single checkpoint pathway. Tumor cells, however, have defects in pathways controlling cell cycle progression such that the perturbation of additional  
10 checkpoints renders them particularly sensitive to DNA damaging agents. For example, tumor cells that contain mutant p53 are defective both in the G1 DNA damage checkpoint and in the ability to maintain the G2 DNA damage checkpoint (Bunz et al., *Science*, 282:1497-1501, 1998).  
15 Checkpoint inhibitors that target initiation of the G2 checkpoint or the S phase checkpoint are expected to further cripple the ability of these tumor cells to repair DNA damage and, therefore, are candidates to enhance the therapeutic index of both radiation and systemic  
20 chemotherapy (Gesner, T., Abstract at SRI Conference: Protein Phosphorylation and Drug Discovery World Summit, March 2003).

In the presence of DNA damage or any impediment to DNA replication, the checkpoint proteins Atm and  
25 Atr initiate a signal transduction pathway leading to cell cycle arrest. Atm has been shown to play a role in a DNA damage checkpoint in response to ionizing radiation. Atr is stimulated by agents that cause double strand DNA breaks, single strand DNA breaks, and agents  
30 that block DNA replication.

Chk1 is a protein kinase that lies downstream from Atm and/or Atr in the DNA damage checkpoint signal transduction pathway (Sanchez et al., *Science*, 277:1497-1501, 1997; U.S. Patent No. 6,218,109). In mammalian

- 7 -

cells, Chk1 is phosphorylated in response to agents that cause DNA damage including ionizing radiation, ultra-violet (UV) light, and hydroxyurea (Sanchez et al., supra; Lui et al., *Genes Dev.*, 14:1448-1459, 2000). This  
5 phosphorylation, which activates Chk1 in mammalian cells, is dependent on Atm (Chen et al., *Oncogene*, 18:249-256, 1999) and Atr (Lui et al., supra). Furthermore, Chk1 has been shown to phosphorylate both weel (O'Connell et al., *EMBO J.*, 16:545-554, 1997) and Pds1 (Sanchez et al.,  
10 *Science*, 286:1166-1171, 1999), gene products known to be important in cell cycle control.

These studies demonstrate that mammalian Chk1 plays a role in the Atm dependent DNA damage checkpoint leading to arrest at S phase. A role for Chk1 in the S  
15 phase mammalian cells has recently been elucidated (Feijoo et al., *J. Cell. Biol.*, 154:913-923, 2001; Zhao et al., *PNAS USA*, 99:14795-800, 2002; Xiao et al., *J Biol Chem.*, 278(24):21767-21773, 2003; Sorensen et al., *Cancer Cell*, 3(3):247-58, 2003) highlighting the role of Chk1 in  
20 monitoring the integrity of DNA synthesis. Chk1 invokes an S-phase arrest by phosphorylating Cdc25A, which regulates cyclinA/cdk2 activity (Xiao et al., supra and Sorensen et al., supra). Chk1 also invokes a G2 arrest by phosphorylating and inactivating Cdc25C, the dual  
25 specificity phosphatase that normally dephosphorylates cyclin-B/cdc2 (also known as Cdk1) as cells progress from G2 into mitosis (Fernery et al., *Science*, 277:1495-7, 1997; Sanchez et al., supra; Matsuoka et al., *Science*, 282:1893-1897, 1998; and Blasina et al., *Curr. Biol.*,  
30 9:1-10, 1999). In both cases, regulation of Cdk activity induces a cell cycle arrest to prevent cells from entering mitosis in the presence of DNA damage or unreplicated DNA.



- 8 -

Additional classes of cell cycle checkpoint inhibitors operate at either the G1 or G2/M phase. UCN-01, or 7-hydroxystaurosporine, originally was isolated as a nonspecific kinase inhibitor having its primary effect on protein kinase C, but recently it has been found to inhibit the activity of Chk1 and abrogate the G2 cell cycle checkpoint (Shi et al., supra). Thus, UCN-01 is a nonselective Chk1 inhibitor, and is toxic to cells at high doses. At low doses, it nonspecifically inhibits many cellular kinases and also inhibits the G1 checkpoint (Tenzer and Pruschy, *Curr. Med. Chem. Anti-Cancer Agents*, 3:35-46, 2003).

UCN-01 has been used in conjunction with cancer therapies, such as radiation, the anti-cancer agent camptothecin (Tenzer and Pruschy, supra), and gemcitabine (Shi et al., supra), with limited success. In addition, UCN-01 also has been used to potentiate the effects of temozolomide (TMZ)-induced DNA mismatch repair (MMR) in glioblastoma cells (Hirose et al., *Cancer Res.*, 61:5843-5849, 2001). In the clinic, UCN-01 is not an effective chemotherapeutic as expected, possibly due to a failure in treatment scheduling and a lack of identification of particular key molecular targets (Grant and Roberts, *Drug Resistance Updates*, 6:15-26, 2003). Thus, Mack et al. report cell cycle-dependent potentiation of cis-platin by UCN-01 in a cultured nonsmall-cell lung carcinoma cell line, but do not identify with specificity the key cell cycle checkpoint(s) targeted by UCN-01. (Mack et al., *Cancer Chemother. Pharmacol.*, 51(4):337-348, 2003).

Several other strategies exist for sensitizing tumor cells to treatment with cell cycle affecting chemotherapeutics. For example, administration of 2-aminopurine abrogates multiple cell cycle checkpoint mechanisms, such as mimosine-induced G1 arrest or hydroxyurea-

- 9 -

induced S phase arrest, allowing the cell to progress into and through mitosis (Andreassen et al., *Proc Natl Acad Sci USA*, 86:2272-2276, 1992). Caffeine, a methylxanthine, has also been used to enhance cytotoxicity of DNA-damaging agents, such as cis-platin and ionizing radiation, by mediating progression through the G2 checkpoint and thereby inducing cell death. (Bracey et al., *Clin. Cancer Res.*, 3:1371-1381, 1997). However, the dose of caffeine used to accomplish the cell cycle abrogation exceeds clinically acceptable levels and is not a viable therapeutic option. Additionally, antisense nucleotides to Chk1 kinase have been used to increase sensitivity to the topoisomerase inhibitor BNP1350 (Yin et al., *Biochem. Biophys. Res. Commun.*, 295:435-44, 2002), but demonstrate problems typically associated with antisense treatment and gene therapy.

Chk1 inhibitors have been disclosed in WO 02/070494, WO 04/014876, and WO 03/101444. Additional Chk1 inhibitors include diarylurea compounds, e.g., aryl- and heteroaryl-substituted urea compounds disclosed in U.S. Patent Publication No. 2003-0069284 A1; methylxanthines and related compounds (Fan et al., *Cancer Res.*, 55:1649-54 (1995); ureidothiophenes (WO 03/029241); N-pyrrolopyridinyl carboxamides (WO 03/28724); antisense Chk1 nucleotides (WO 01/57206); Chk1 receptor antagonists (WO 00/16781); heteroaromatic carboxamide derivatives (WO 03/037886); aminothiophenes (WO 03/029242); (indazolyl)-benzimidazoles (WO 03/004488); heterocyclic-hydroxyimino-fluorenes (WO 02/16326); scytoneman skeleton-containing derivatives (scytonemin) (U.S. Patent No. 6,495,586); heteroarylbenzamides (WO 01/53274); indazole compounds (WO 01/53268); indolacarbazoles (see Tenzer et al., *supra*); chromane derivatives (WO 02/070515); paullones (Schultz et al., *J. Med. Chem.*, 42:2909-2919 (1999));

- 10 -

indenopyrazoles (WO 99/17769); flavones (Sedlacek et al.,  
Int. J. Oncol., 9:1143-1168 (1996); peptide derivatives  
of peptide loop of serine threonine kinases (WO  
98/53050); and oxindoles (WO 03/051838).


5                   However, a need still exists in the art for  
effective and selective inhibitors of Chk1. The present  
invention addresses this and other needs.

#### SUMMARY OF THE INVENTION

The present invention relates to potent and  
10   selective inhibitors of the checkpoint kinase Chk1. The  
present Chk1 inhibitors are useful in treating indica-  
tions involving aberrant cell proliferation, and as  
chemosensitizing and radiosensitizing agents in the  
treatment of indications related to DNA damage or lesions  
15   in DNA replication.

Therefore, one aspect of the present invention  
is to provide compounds of structural formula (I). The  
compounds are useful in a method of inhibiting Chk1 com-  
prising a step of administering an effective amount of a  
20   compound of structural formula (I) to an individual.

Compounds of formula (I) have a structural  
formula:



(I),

wherein X<sup>1</sup> is null, -O-, -S-, -CH<sub>2</sub>-, or  
25   -N(R<sup>1</sup>)-;  
X<sup>2</sup> is -O-, -S-, or -N(R<sup>1</sup>)-;

- 11 -

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, and C<sub>1-6</sub>alkyl substituted with a hetero-  
5 aryl or aryl group, wherein (a) said aryl or heteroaryl group of W is substituted with at least one of CF<sub>3</sub> and heteroaryl, (b) said aryl group of group W is optionally substituted with one to three substituents represented by R<sup>2</sup>, and (c) said heteroaryl group of group W is optional-  
10 ly substituted with one to three substituents represented by R<sup>5</sup>;

R<sup>1</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, and aryl;

R<sup>2</sup> is selected from the group consisting of  
15 heteroaryl, halo, optionally substituted C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, OCF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N-(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyl-eneC(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyl-eneOR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyl-  
20 eneSO<sub>2</sub>NR<sup>3</sup>, C<sub>1-6</sub>alkyleneOR<sup>3</sup>, and SR<sup>3</sup>;


R<sup>3</sup> is selected from the group consisting of hydro, halo, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, cycloalkyl, aryl, heteroaryl, CO<sub>2</sub>R<sup>4</sup>, SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocyclo-  
25 alkyl, N(R<sup>4</sup>)<sub>2</sub>, and SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkylenearyl, C<sub>1-6</sub>alkylenehet-eroaryl, C<sub>1-6</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl, C<sub>1-6</sub>alkylene-SO<sub>2</sub>aryl, optionally substituted C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>2</sub>, OCF<sub>3</sub>, C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>3</sub><sup>+</sup>, C<sub>3-8</sub>heterocycloalkyl, and CH(C<sub>1-6</sub>alkyl-eneN(R<sup>4</sup>)<sub>2</sub>)<sub>2</sub>, or two R<sup>3</sup> groups are taken together to form  
30 an optionally substituted 3- to 6-membered aliphatic ring;

R<sup>4</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, cycloalkyl, aryl, heteroaryl, C<sub>1-6</sub>alkyl-enearyl, and SO<sub>2</sub>C<sub>1-6</sub>alkyl, or two R<sup>4</sup> groups are taken

- 12 -

together to form an optionally substituted 3- to 6-membered ring;

$R^5$  is selected from the group consisting of  $C_{1-6}$ alkyl, aryl, heteroaryl, heterocycloalkyl,  $N(R^3)_2$ ,  $OR^3$ ,  
 5 halo,  $N_3$ , CN,  $C_{1-6}$ alkylenearyl,  $C_{1-6}$ alkylene $N(R^3)_2$ ,  $C(O)R^3$ ,  
 $C(O)OR^3$ ,  $C(O)N(R^3)_2$ ,  $N(R^1)C(O)R^3$ ,  $N(R^1)C(O)OR^3$ ,  $CF_3$ , and



$R^6$  is selected from the group consisting of  $OR^{11}$ ,  $-C\equiv C-R^7$ , and heteroaryl;

10  $R^7$  is selected from the group consisting of hydro,  $C_{1-6}$ alkyl, aryl,  $C_{1-6}$ alkylenearyl, heteroaryl,  $C_{1-6}$ alkyleneheteroaryl, alkoxy;

$R^8$ ,  $R^9$ , and  $R^{10}$ , independently, are selected from the group consisting of hydro, halo, optionally  
 15 substituted  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $OCF_3$ ,  $CF_3$ ,  $NO_2$ , CN, NC,  $N(R^3)_2$ ,  $OR^3$ ,  $CO_2R^3$ ,  $C(O)N(R^3)_2$ ,  $C(O)R^3$ ,  $N(R^1)COR^3$ ,  $N(R^1)C(O)OR^3$ ,  $N(R^8)C(O)OR^3$ ,  $N(R^1)C(O)C_{1-6}$ alkylene-  
 $C(O)R^3$ ,  $N(R^1)C(O)C_{1-6}$ alkylene $C(O)OR^3$ ,  $N(R^1)C(O)C_{1-6}$ alkylene-  
 $OR^3$ ,  $N(R^1)C(O)C_{1-6}$ alkyleneNHC(O) $OR^3$ ,  $N(R^1)C(O)C_{1-6}$ alkylene-  
 20  $SO_2NR^3$ ,  $C_{1-6}$ alkylene $OR^3$ , and  $SR^3$ ;

$R^{11}$  is selected from the group consisting of hydro,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl, cycloalkyl, heterocyclo-  
 alkyl, aryl, heteroaryl,  $SO_2R^4$ ,  $C_{1-6}$ alkyl substituted with  
 one or more of halo, hydroxy, aryl, heteroaryl,  $N(R^4)_2$ ,  
 25 and  $SO_2R^4$ ,  $C_{1-6}$ alkylenearyl,  $C_{1-6}$ alkyleneheteroaryl,  
 $C_{1-6}$ alkylene $C_{3-8}$ heterocycloalkyl,  $C_{1-6}$ alkylene $SO_2$ aryl,  
 optionally substituted  $C_{1-6}$ alkylene $N(R^4)_2$ ,  $OCF_3$ ,  $C_{1-6}$ alkyl-

- 13 -

eneN(R<sup>4</sup>)<sub>3</sub>\*, C<sub>3-8</sub>heterocycloalkyl, and CH(C<sub>1-6</sub>alkylene-N(R<sup>4</sup>)<sub>2</sub>)<sub>2</sub>;

or a pharmaceutically acceptable salt, or prodrug, or solvate thereof.

5           Another aspect of the present invention is to provide pharmaceutical compositions comprising one or more compound of structural formula (I), and use of the compositions in a therapeutic treatment of a disease or disorder, wherein inhibition of Chk1, *in vivo* or *ex vivo*,  
10           provides a therapeutic benefit or is of research or diagnostic interest.


          Yet another aspect of the present invention is to provide a method of sensitizing cells in an individual undergoing a chemotherapeutic or radiotherapeutic treat-  
15           ment for a medical condition comprising administration of a compound of structural formula (I) in combination with a chemotherapeutic agent, a radiotherapeutic agent, or both, to the individual. A nonlimiting indication treated by this method is a cancer.

20           Another aspect of the present invention is to provide a method of inhibiting or preventing aberrant cell proliferation. In one embodiment, a method comprises contacting a cell population comprising aberrantly proliferating cells with at least one Chk1 activator in  
25           an amount and for a time sufficient to substantially synchronize cell cycle arrest among the aberrantly proliferating cells. Upon achieving substantial synchronization of cell cycle arrest in the cell population, the cell population is contacted with at least one Chk1  
30           inhibitor in an amount and for a time sufficient to substantially abrogate the cell cycle arrest.

          These and other aspects of the present invention will become apparent from the following detailed description of the preferred embodiments.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Compounds of the present invention have a structural formula (I):



5                    wherein  $X^1$  is null, -O-, -S-, -CH<sub>2</sub>-, or -N(R<sup>1</sup>)-;

$X^2$  is -O-, -S-, or -N(R<sup>1</sup>)-;

$Y$  is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

10                     $W$  is selected from the group consisting of heteroaryl, aryl, and C<sub>1-6</sub>alkyl substituted with a heteroaryl or aryl group, wherein (a) said aryl or heteroaryl group of group  $W$  is substituted with at least one of CF<sub>3</sub> and heteroaryl; (b) said aryl group of group  $W$  is  
15 optionally substituted with one to three substituents represented by R<sup>2</sup>, and (c) said heteroaryl group of group  $W$  is optionally substituted with one to three substituents represented by R<sup>5</sup>;

20                    R<sup>1</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, and aryl;


                    R<sup>2</sup> is selected from the group consisting of heteroaryl, halo, optionally substituted C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, OCF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)-  
25 C<sub>1-6</sub>alkyleneC(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneOR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneSO<sub>2</sub>NR<sup>3</sup>, C<sub>1-6</sub>alkyleneOR<sup>3</sup>, and SR<sup>3</sup>;

- 15 -

$R^3$  is selected from the group consisting of hydro, halo,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl, cycloalkyl, aryl, heteroaryl,  $CO_2R^4$ ,  $SO_2R^4$ ,  $C_{1-6}$ alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocyclo-  
 5 alkyl,  $N(R^4)_2$ , and  $SO_2R^4$ ,  $C_{1-6}$ alkylenearyl,  $C_{1-6}$ alkyleneheteroaryl,  $C_{1-6}$ alkylene $C_{3-8}$ heterocycloalkyl,  $C_{1-6}$ alkylene- $SO_2$ aryl, optionally substituted  $C_{1-6}$ alkylene $N(R^4)_2$ ,  $OCF_3$ ,  $C_{1-6}$ alkylene $N(R^4)_3^+$ ,  $C_{3-8}$ heterocycloalkyl, and  $CH(C_{1-6}$ alkylene $N(R^4)_2)_2$ , or two  $R^3$  groups are taken together to form  
 10 an optionally substituted 3- to 6-membered aliphatic ring;

$R^4$  is selected from the group consisting of hydro,  $C_{1-6}$ alkyl, cycloalkyl, aryl, heteroaryl,  $C_{1-6}$ alkylenearyl, and  $SO_2C_{1-6}$ alkyl, or two  $R^4$  groups are taken  
 15 together to form an optionally substituted 3- to 6-membered ring;

$R^5$  is selected from the group consisting of  $C_{1-6}$ alkyl, aryl, heteroaryl, heterocycloalkyl,  $N(R^3)_2$ ,  $OR^3$ , halo,  $N_3$ , CN,  $C_{1-6}$ alkylenearyl,  $C_{1-6}$ alkylene $N(R^3)_2$ ,  $C(O)R^3$ ,  
 20  $C(O)OR^3$ ,  $C(O)N(R^3)_2$ ,  $N(R^1)C(O)R^3$ ,  $N(R^1)C(O)OR^3$ ,  $CF_3$ , and



$R^6$  is selected from the group consisting of  $OR^{11}$ ,  $-C\equiv C-R^7$ , and heteroaryl;

$R^7$  is selected from the group consisting of  
 25 hydro,  $C_{1-6}$ alkyl, aryl,  $C_{1-6}$ alkylenearyl, heteroaryl,  $C_{1-6}$ alkyleneheteroaryl, alkoxy;

$R^8$ ,  $R^9$ , and  $R^{10}$ , independently, are selected from the group consisting of hydro, halo, optionally



- 16 -

substituted C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, OCF<sub>3</sub>, CF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>8</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneOR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneSO<sub>2</sub>NR<sup>3</sup>, C<sub>1-6</sub>alkyleneOR<sup>3</sup>, and SR<sup>3</sup>;

R<sup>11</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, N(R<sup>4</sup>)<sub>2</sub>, and SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkylenearyl, C<sub>1-6</sub>alkyleneheteroaryl, C<sub>1-6</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl, C<sub>1-6</sub>alkyleneSO<sub>2</sub>aryl, optionally substituted C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>2</sub>, OCF<sub>3</sub>, C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>3</sub><sup>+</sup>, C<sub>3-8</sub>heterocycloalkyl, and CH(C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>2</sub>)<sub>2</sub>;

and a pharmaceutically acceptable salt, or prodrug, or solvate thereof.

Preferred compounds of the present invention are those wherein X<sup>1</sup> and X<sup>2</sup> are -N(H)-;







Y is O or S; and

Preferably W is heteroaryl. In one embodiment, W is heteroaryl containing at least two heteroatoms selected from the group consisting of N, O, and S, said heteroaryl ring optionally substituted with one or two substituents selected from the group consisting of optionally substituted C<sub>1-6</sub>alkyl, aryl, heteroaryl, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, CO<sub>2</sub>R<sup>3</sup>, CN, and halo, wherein R<sup>3</sup> is as previously defined.


Other preferred compounds of structural formula (I) are those wherein W is selected from the group consisting of pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, optionally substituted with one or two substituents selected from the group consisting of C<sub>1-6</sub>alkyl, aryl, heteroaryl, N(R<sup>3</sup>)<sub>2</sub>, C(O)N(R<sup>3</sup>)<sub>2</sub>, CO<sub>2</sub>R<sup>3</sup>, OR<sup>3</sup>, and halo.

- 17 -

In some preferred embodiments, W is selected from the group consisting of



, and




optionally substituted with one to four substituents selected from the group consisting of C<sub>1-6</sub>alkyl, C<sub>2-6</sub>.


- 18 -

alkynyl, aryl, heteroaryl, CN, CO<sub>2</sub>R<sup>3</sup>, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, and halo.

In more preferred embodiments, W is



or






5









In a most preferred embodiment, W is pyrazinyl and X<sup>1</sup> and X<sup>2</sup> each are N(H).

In yet another preferred embodiment, the heteroaryl group substituent on W and the heteroaryl group of R<sup>6</sup>, independently, are selected from the group consisting of










10











- 19 -




- 20 -




- 21 -





, and



As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight chain and branched propyl and butyl groups. Unless otherwise indicated, the hydrocarbon group can contain up to 20 carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a C<sub>6</sub>-C<sub>16</sub> bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. Alkyl groups optionally can be substituted, for example, with hydroxy (OH), halo, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, amino (N(R<sup>3</sup>)<sub>2</sub>), and sulfonyl (SO<sub>2</sub>R<sup>3</sup>), wherein R<sup>3</sup> is as previously defined.

The term "cycloalkyl" is defined as a cyclic C<sub>3-8</sub> hydrocarbon group, e.g., cyclopropyl, cyclobutyl, cyclohexyl, or cyclopentyl. "Heterocycloalkyl" is defined similarly as cycloalkyl, except the ring contains one to three heteroatoms independently selected from the group consisting of oxygen, nitrogen, and sulfur. Cycloalkyl and heterocycloalkyl groups can be saturated or partially unsaturated ring systems optionally substituted with, for example, one to three groups, independently selected from the group consisting of C<sub>1-4</sub>alkyl, C<sub>1-3</sub>alkyl-eneOH, C(O)NH<sub>2</sub>, NH<sub>2</sub>, oxo (=O), aryl, trifluoroethanoyl, and OH. Heterocycloalkyl groups optionally can be

- 23 -

further N-substituted with C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, C<sub>1-3</sub>alkylenearyl, or C<sub>1-3</sub>alkyleneheteroaryl.

The term "alkenyl" is defined identically as "alkyl," except the group contains a carbon-carbon double  
5 bond.

The term "alkynyl" is defined identically as "alkyl," except the group contains a carbon-carbon triple bond.

The term "alkylene" refers to an alkyl group  
10 having a substituent. For example, the term "C<sub>1-6</sub>alkyleneC(O)OR" refers to an alkyl group containing one to six carbon atoms substituted with a -C(O)OR group. The alkylene group is optionally substituted with one or more substituent previously listed as an optional alkyl substituent.  
15

The term "halo" or "halogen" is defined herein as fluorine, bromine, chlorine, and iodine.

The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic  
20 group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl. Unless otherwise indicated, an aryl group can be unsubstituted or substituted with one or more, and in particular one to four groups independently selected from, for example, halo, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, OCF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N-(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkyleneC(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkyleneOR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkyleneSO<sub>2</sub>NR<sup>3</sup>, C<sub>1-3</sub>alkyleneOR<sup>1</sup>, and SR<sup>3</sup>,  
25 wherein R<sup>1</sup> and R<sup>3</sup> are as previously defined. Exemplary aryl groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl, chlorophenyl, methylphenyl, methoxyphenyl, trifluoromethylphenyl, nitrophenyl, 2,4-methoxychlorophenyl, and the like. The terms "aryl-

30



- 24 -

C<sub>1-3</sub>alkyl" and "heteroarylC<sub>1-3</sub>alkyl" are defined as an aryl or heteroaryl group having a C<sub>1-3</sub>alkyl substituent.

The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one or two  
5 aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring. Unless otherwise indicated, a heteroaryl group can be unsubstituted or substituted with one or more, and in particular one to four, substituents selected from, for  
10 example, C<sub>1-6</sub>alkyl, aryl, heteroaryl, CF<sub>3</sub>, CN, C(O)N(R<sup>3</sup>)<sub>2</sub>, CO<sub>2</sub>R<sup>2</sup>, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, and halo, wherein R<sup>3</sup> is as previously defined. Examples of heteroaryl groups include, but are not limited to, thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazinyl, triazolyl,  
15 isothiazolyl, isoxazolyl, imidazolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "hydro" is defined as -H.

The term "hydroxy" is defined as -OH.

The term "nitro" is defined as -NO<sub>2</sub>.

20 The term "cyano" is defined as -CN.

The term "isocyano" is defined as -NC.

The term "trifluoromethoxy" is defined as -OCF<sub>3</sub>.

The term "azido" is defined as -N<sub>3</sub>.


25 The term "3- to 8-membered ring" as used herein refers to carbocyclic and heterocyclic aliphatic or aromatic groups, including, but not limited to, morpholinyl, piperidinyl, phenyl, thiophenyl, furyl, pyrrolyl, imidazolyl, pyrimidinyl, and pyridinyl, op-  
30 tionally substituted with one or more, and in particular one to three, groups exemplified above for aryl groups.

The carbon atom content of hydrocarbon-containing moieties is indicated by a subscript designating the minimum and maximum number of carbon atoms in the


moiety, e.g., "C<sub>1-6</sub>alkyl" refers to an alkyl group having one to six carbon atoms, inclusive.

In the structures herein, for a bond lacking a substituent, the substituent is methyl, for example,

5



When no substituent is indicated as attached to a carbon atom on a ring, it is understood that the carbon atom contains the appropriate number of hydrogen atoms. In addition, when no substituent is indicated as attached to a carbonyl group or a nitrogen atom, for example, the substituent is understood to be hydrogen, e.g.,



The abbreviation "Me" is methyl. The abbreviation CO and C(O) is carbonyl (C=O).

The notation N(R<sup>x</sup>) (wherein x represents an alpha or numeric character, such as for example R<sup>a</sup>, R<sup>b</sup>, R<sup>3</sup>, R<sup>4</sup>, and the like) is used to denote two R<sup>x</sup> groups attached to a common nitrogen atom. When used in such notation, the R<sup>x</sup> group can be the same or different, and are selected from the group as defined by the R<sup>x</sup> group.

"Chk1 inhibitor" means any compound, known or after-discovered whether naturally occurring or synthetic, that is capable of at least partially abrogating cell cycle checkpoint activity of the Chk1 protein. Abrogation of cell cycle checkpoint is achieved when the

cellular checkpoint mechanism(s) is (are) overcome sufficiently to allow the cell to pass from the cell cycle phase in which it is halted to the next phase in the cell cycle or to allow the cell to pass directly to cell

5 death. Abrogation of the cell cycle checkpoint permits cells to carry damage or imperfections to subsequent cell cycle phases, thereby inducing or promoting cell death. Cell death can occur by any associated mechanism, including apoptosis and mitotic catastrophe.

10 "Chk1 activator" means any known or after-discovered agent having the ability to activate Chk1 kinase activity in DNA repair and homeostasis at cell cycle checkpoints, and thus induce at least partial cell cycle arrest. Chk1 activators include agents capable of  
15 arresting the cell cycle at any phase of the cell cycle, which phase may be referred to herein as the "target phase" for that activator. Target phases include any of the cell cycle phases except mitosis, i.e., the G1 phase, S phase, and G2 phase. Chk1 activators useful in the  
20 invention include DNA damaging agents, such as chemotherapeutic agents and/or radiation (or "radiotherapeutic agents"), such as ionizing or ultraviolet radiation. Radiation Chk1 activators include, but are not limited to, gamma radiation, X-ray radiation, ultraviolet light,  
25 visible light, infrared radiation, microwave radiation, and mixtures thereof.

"Inhibiting aberrant cell proliferation" means to retard or eliminate the rate at which aberrantly proliferating cells proliferate. This inhibition can result  
30 either from a decreased rate of replication, an increased rate of cell death, or both. Cell death can occur by any mechanism, including apoptosis and mitotic catastrophe.

"Preventing aberrant cell proliferation" means inhibiting aberrant cell proliferation prior to occurrence, or inhibiting the recurrence thereof.

"In vivo" means within a living subject, as  
5 within an animal or human. In this context, agents can be used therapeutically in a subject to retard or eliminate the proliferation of aberrantly replicating cells. The agents also can be used as a prophylactic to prevent the occurrence or recurrence of aberrant cell proliferation or the manifestation of symptoms associated there-  
10 with.

"Ex vivo" means outside a living subject. Examples of *ex vivo* cell populations include *in vitro* cell cultures and biological samples such as fluid or  
15 tissue samples from humans or animals. Such samples can be obtained by methods well known in the art. Exemplary biological fluid samples include blood, cerebrospinal fluid, urine, saliva. Exemplary tissue samples include tumors and biopsies thereof. In this context, the  
20 present compounds can be in numerous applications, both therapeutic and experimental.

The term "radiosensitizer," as used herein, is defined as a compound, administered to a human or other animal in a therapeutically effective amount to increase  
25 the sensitivity of cells to electromagnetic radiation and/or to promote the treatment of diseases treatable with electromagnetic radiation.

The terms "electromagnetic radiation" and "radiation" as used herein include, but are not limited  
30 to, radiation having the wavelength of 10-20 to 100 nanometers.

The present invention includes all possible stereoisomers and geometric isomers of the compounds of structural formula (I). The present invention includes

- 28 -

not only racemic compounds, but optically active isomers as well. When a compound of structural formula (I) is desired as a single enantiomer, it can be obtained either by resolution of the final product or by stereospecific  
5 synthesis from either isomerically pure starting material or use of a chiral auxiliary reagent, for example, see Ma et al., *Tetrahedron: Asymmetry*, 8(6), pages 883-888 (1997). Resolution of the final product, an intermediate, or a starting material can be achieved by any suitable method known in the art. Additionally, in situations where tautomers of the compounds of structural  
10 formula (I) are possible, the present invention is intended to include all tautomeric forms of the compounds. As demonstrated hereafter, specific stereoisomers can  
15 exhibit an exceptional ability to inhibit Chk1 in combination with chemotherapeutic or radiotherapeutic treatments.

Prodrugs of compounds of structural formula (I) also can be used as the compound in a method of the  
20 present invention. It is well established that a prodrug approach, wherein a compound is derivatized into a form suitable for formulation and/or administration, then released as a drug *in vivo*, has been successfully employed to transiently (e.g., bioreversibly) alter the physico-  
25 chemical properties of the compound (see, H. Bundgaard, Ed., "Design of Prodrugs," Elsevier, Amsterdam, (1985); Silverman, "The Organic Chemistry of Drug Design and Drug Action," Academic Press, San Diego, chapter 8, (1992); Hillgren et al., *Med. Res. Rev.*, 15, 83 (1995)).

30 Compounds of the present invention can contain one or more functional groups. The functional groups, if desired or necessary, can be modified to provide a prodrug. Suitable prodrugs include, for example, acid derivatives, such as amides and esters. It also is

- 29 -

appreciated by those skilled in the art that N-oxides can be used as prodrugs.

As used herein, the term "pharmaceutically acceptable salts" refers compounds of structural formula (I) that contain an acidic moiety and form salts having suitable cations. Suitable pharmaceutically acceptable cations include alkali metal (e.g., sodium or potassium) and alkaline earth metal (e.g., calcium or magnesium) cations. In addition, the pharmaceutically acceptable salts of compounds of structural formula (I) that contain a basic center are acid addition salts formed with pharmaceutically acceptable acids. Nonlimiting examples include the hydrochloride, hydrobromide, sulfate, bisulfate, phosphate, hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzene sulfonate, and p-toluenesulphonate salts. In light of the foregoing, any reference to compounds of the present invention appearing herein is intended to include compounds of structural formula (I) as well as pharmaceutically acceptable salts or solvates thereof.

The compounds of the present invention can be therapeutically administered as the neat chemical, but it may be useful to administer compounds of structural formula (I) as a pharmaceutical composition or formulation. Thus, the present invention provides a pharmaceutical composition comprising a compound of the formula (I) together with a pharmaceutically acceptable diluent or carrier therefor. Also provided is a process of preparing a pharmaceutical composition comprising admixing a compound of formula (I) with a pharmaceutically acceptable diluent or carrier therefor.

Accordingly, the present invention further provides pharmaceutical formulations comprising a com-

pound of structural formula (I), or a pharmaceutically acceptable salt, prodrug, or solvate thereof, together with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic and/or prophylactic ingredients. The carriers are "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Such carriers may be found for example, in *Remington's Pharmaceutical Sciences*, 17<sup>th</sup> Ed., Mack Publishing Co., Easton, PA. (1985).

Inhibition of the checkpoint kinase typically is measured using a dose-response assay in which a sensitive assay system is contacted with a compound of interest over a range of concentrations, including concentrations at which no or minimal effect is observed, through higher concentrations at which partial effect is observed, to saturating concentrations at which a maximum effect is observed. Theoretically, such assays of the dose-response effect of inhibitor compounds can be described as a sigmoidal curve expressing a degree of inhibition as a function of concentration. The curve also theoretically passes through a point at which the concentration is sufficient to reduce activity of the checkpoint enzyme to a level that is 50% that of the difference between minimal and maximal enzyme activity in the assay. This concentration is defined as the "Inhibitory Concentration (50%)" or "IC<sub>50</sub>" value. Determination of IC<sub>50</sub> values may be made using conventional biochemical (acellular) assay techniques or cell-based assay techniques.

Comparisons of the efficacy of inhibitors often are provided with reference to comparative IC<sub>50</sub> values, wherein a higher IC<sub>50</sub> indicates that the test compound is less potent, and a lower IC<sub>50</sub> indicates that

the compound is more potent, than a reference compound. Compounds of the present invention demonstrate an  $IC_{50}$  value of less than 5  $\mu$ M, and down to 0.1 nM, when measured using the dose-response assay. Preferred compounds demonstrate an  $IC_{50}$  value of 500 nM or less. More preferred compounds of the present invention demonstrate an  $IC_{50}$  value of less than 250 nM or less, 100 nM or less, 50 nM or less, 20 nM or less.

Preferred Chk1 inhibitors of the invention are selective, i.e., demonstrate at least a 20-fold selectivity in inhibiting Chk1 over the following protein kinases: protein kinase A, protein kinase C, cdc2, and pp60v-src. More preferred Chk1 inhibitors of the present invention preferably exhibit at least 75-fold selectivity in inhibiting Chk1 over the following protein kinases: protein kinase A, protein kinase C, cdc2, and pp60v-src. Most preferred Chk1 inhibitors of the present invention demonstrate at least 100-fold selectivity against protein kinase A, protein kinase C, cdc2, pp60v-src, protein kinase B/Akt-1, p38MapK, ERK1, p70S6K, cdc2, cdk2, Chk2, and the abl tyrosine kinase. "Fold selectivity" is defined as the  $IC_{50}$  of the Chk1 inhibitor for the comparison kinase divided by the  $IC_{50}$  of the Chk1 inhibitor for Chk1.

Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effective amount" means an amount sufficient to treat an individual suffering an indication, or to alleviate the existing symptoms of the indication. Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.



- 32 -

In addition to the Chk1 inhibitor, pharmaceutical compositions of the invention can be formulated to include cytokines, lymphokines, growth factors, other hematopoietic factors, or mixtures thereof, to reduce  
5 adverse side effects that can arise from, or be associated with, administration of the pharmaceutical composition alone. Cytokines, lymphokines, growth factors, or other hematopoietic factors particularly useful in pharmaceutical compositions of the invention include, but are  
10 not limited to, M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IFN, TNF, G-CSF, Meg-CSF, GM-CSF, thrombopoietin, stem cell factor, erythropoietin, angiopoietins, including Ang-1, Ang-2,  
15 Ang-4, Ang-Y, and/or the human angiopoietin-like polypeptide, vascular endothelial growth factor (VEGF), angiogenin, bone morphogenic protein-1 (BMP-1), BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9, BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP receptor IA, BMP receptor IB, brain derived neurotrophic factor,  
20 ciliary neurotrophic factor, ciliary neurotrophic factor receptor cytokine-induced neutrophil chemotactic factor 1, cytokine-induced neutrophil chemotactic factor 2, cytokine-induced neutrophil chemotactic factor 2, endothelial cell growth factor, endothelin 1, epidermal growth factor, epithelial-derived neutrophil attractant, fibroblast growth factor (FGF) 4, FGF 5, FGF 6, FGF 7, FGF 8, FGF 8b, FGF 8c, FGF 9, FGF 10, FGF acidic, FGF basic, glial cell line-derived neurotrophic factor receptor  
25 1, glial cell line-derived neurotrophic factor receptor 2, growth related protein, growth related protein, growth related protein, growth related protein, heparin binding epidermal growth factor, hepatocyte growth factor, hepatocyte growth factor receptor, insulin-like growth factor  
30

- 33 -

I, insulin-like growth factor receptor, insulin-like growth factor II, insulin-like growth factor binding protein, keratinocyte growth factor, leukemia inhibitory factor, leukemia inhibitory factor receptor, nerve growth factor nerve growth factor receptor, neurotrophin-3, neurotrophin-4, placenta growth factor, placenta growth factor 2, platelet-derived endothelial cell growth factor, platelet derived growth factor, platelet derived growth factor A chain, platelet derived growth factor AA, platelet derived growth factor AB, platelet derived growth factor B chain, platelet derived growth factor BB, platelet derived growth factor receptor, platelet derived growth factor receptor, pre-B cell growth stimulating factor, stem cell factor, stem cell factor receptor, transforming growth factor (TGF), TGF, TGF 1, TGF 1.2, TGF 2, TGF 3, TGF 5, latent TGF 1, TGF, binding protein I, TGF binding protein II, TGF binding protein III, tumor necrosis factor receptor type I, tumor necrosis factor receptor type II, urokinase-type plasminogen activator receptor, vascular endothelial growth factor, and chimeric proteins and biologically or immunologically active fragments thereof.

The compounds of structural formula (I) also can be conjugated or linked to auxiliary moieties that promote a beneficial property of the compound in a method of therapeutic use. Such conjugates can enhance delivery of the compounds to a particular anatomical site or region of interest (e.g., a tumor), enable sustained therapeutic concentrations of the compounds in target cells, alter pharmacokinetic and pharmacodynamic properties of the compounds, and/or improve the therapeutic index or safety profile of the compounds. Suitable auxiliary moieties include, for example, amino acids, oligopeptides, or polypeptides, e.g., antibodies such as

monoclonal antibodies and other engineered antibodies;  
and natural or synthetic ligands to receptors in target  
cells or tissues. Other suitable auxiliaries include  
fatty acid or lipid moieties that promote biodistribution  
5 and/or uptake of the compound by target cells (see, e.g.,  
Bradley et al., *Clin. Cancer Res.* (2001) 7:3229).

Formulations of the present invention can be  
administered in a standard manner for the treatment of  
the indicated diseases, such as by oral, parenteral,  
10 transmucosal (e.g., sublingual or via buccal administra-  
tion), topical, transdermal, rectal, or inhalation (e.g.,  
nasal or deep lung inhalation) administration. Par-  
enteral administration includes, but is not limited to  
intravenous, intraarterial, intraperitoneal, subcutane-  
15 ous, intramuscular, intrathecal, and intraarticular modes  
of administration. Parenteral administration also can be  
accomplished using a high pressure technique, like  
POWDERJECT™.

For oral administration and buccal adminis-  
20 tration, the composition can be in the form of tablets or  
lozenges formulated in conventional manner. For example,  
tablets and capsules for oral administration can contain  
conventional excipients such as binding agents (for  
example, syrup, acacia, gelatin, sorbitol, tragacanth,  
25 mucilage of starch, or polyvinylpyrrolidone), fillers  
(for example, lactose, sugar, microcrystalline cellulose,  
maize-starch, calcium phosphate, or sorbitol), lubricants  
(for example, magnesium stearate, stearic acid, talc,  
polyethylene glycol or silica), disintegrants (for  
30 example, potato starch or sodium starch glycolate), or  
wetting agents (for example, sodium lauryl sulfate). The  
tablets can be coated according to methods well known in  
the art.

- 35 -

Alternatively, compounds of the present invention can be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, for example. Moreover, formulations containing these compounds can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can contain conventional additives, for example suspending agents, such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, hydroxypropyl-methylcellulose, carboxymethylcellulose, aluminum stearate gel, and hydrogenated edible fats; emulsifying agents, such as lecithin, sorbitan monooleate, or acacia; nonaqueous vehicles (which can include edible oils), such as almond oil, fractionated coconut oil, oily esters, propylene glycol, and ethyl alcohol; and preservatives, such as methyl or propyl p-hydroxybenzoate and sorbic acid.

Such preparations also can be formulated as suppositories, e.g., containing conventional suppository bases, such as cocoa butter or other glycerides. Compositions for inhalation typically can be provided in the form of a solution, suspension, or emulsion that can be administered as a dry powder or in the form of an aerosol using a conventional propellant, such as dichlorodifluoromethane or trichlorofluoromethane. Typical topical and transdermal formulations comprise conventional aqueous or nonaqueous vehicles, such as eye drops, creams, ointments, lotions, and pastes, or are in the form of a medicated plaster, patch, or membrane.

Additionally, compositions of the present invention can be formulated for parenteral administration by injection or continuous infusion. Formulations for injection can be in the form of suspensions, solutions,

- 36 -

or emulsions in oily or aqueous vehicles, and can contain formulation agents, such as suspending, stabilizing, and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a  
5 suitable vehicle (e.g., sterile, pyrogen-free water) before use.

A composition of the present invention also can be formulated as a depot preparation. Such long acting formulations can be administered by implantation  
10 (for example, subcutaneously or intramuscularly) or by intramuscular injection. Accordingly, the compounds of the invention can be formulated with suitable polymeric or hydrophobic materials (e.g., an emulsion in an acceptable oil), ion exchange resins, or as sparingly  
15 soluble derivatives (e.g., a sparingly soluble salt).

For veterinary use, a compound of formula (I), or a pharmaceutically acceptable salt, prodrug, or solvent thereof, is administered as a suitably acceptable formulation in accordance with normal veterinary practice.  
20 The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal. Animals treatable by the present compounds and methods include, but are not limited to, pets, livestock, show animals, and zoo specimens.  
25


#### SYNTHETIC METHODS

Compounds of the present invention can be prepared by the following synthetic schemes. Starting materials can be obtained from commercial sources or  
30 prepared by well-established literature methods known to those of ordinary skill in the art. The groups X, R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, and R<sup>11</sup> are defined above unless otherwise noted below. R<sup>2</sup> is as defined above, and also

- 37 -


includes CF<sub>3</sub> and heteroaryl in the following synthetic schemes.

Scheme 1




5                    As illustrated in Scheme 1, compounds of  
 formula 4 can be prepared from compounds of formula 6 by  
 treatment with a base, such as DIEA, and diphenyl  
 phosphoryl azide. A typical solvent for this reaction is  
 THF, and the reaction is performed behind a blast shield  
 10 at room temperature over a one- to twelve-hour period.

Scheme 2




- 38 -



Scheme 2 shows an alternative synthesis of compounds of formula 5. Compounds of formula 3 are treated with compounds of formula 7, which is prepared according to Scheme 3. A useful, nonlimiting solvent is DMF, and the reaction temperature is maintained between room temperature and 60°C for about one to twelve hours.





Scheme 3



As demonstrated in Scheme 3, compounds of formula 7 can be prepared from compounds of formula 8 by treatment with an aryl chloroformate, such as phenyl chloroformate or p-nitrophenyl chloroformate, in the presence of a base, such as pyridine. Nonlimiting solvents used in this reaction include CH<sub>2</sub>Cl<sub>2</sub> or pyridine, at temperatures from 0°C to room temperature.

- 39 -

Scheme 4

 $\text{X}=\text{Otf}, \text{Br}, \text{I}$ 

5

Scheme 4 shows an approach to compounds of formula 3. Compounds of formula 1 are converted to compounds of formula 2 by treatment with aryl boronic




- 40 -

acids and a source of palladium(0) (for example, palladium tetrakis triphenylphosphine) in the presence of a basic aqueous solution, such as sodium bicarbonate, potassium carbonate, or potassium phosphate. Nonlimiting  
5 examples of solvents used in this reaction include THF, dioxane, or ethylene glycol dimethyl ether. The reaction typically is performed at temperatures between 0°C and 90°C for about 1 to 12 hours. Compounds of formula 2 are converted to compounds of formula 3 in the presence of  
10 palladium on carbon, platinum on carbon, or zinc, for example. Examples of solvents used in this reaction include, but are not limited to, methanol, ethanol, or acetic acid. Alternatively, compounds of formula 1 can be used to arylate terminal alkynes, 11, using a cat-  
15 alyst, such as dichloro palladium bis triphenyl phosphine or any other source of palladium(0). Reactions typically are conducted at temperatures varying from room temperature to 90°C, in the presence of a base, such as trimethylamine.


20 Furthermore, compounds of formula 1, where X is a triflate, i.e., tf, can be obtained from compounds of formula 9. Typical reagents include triflic anhydride or N-phenyl triflimide. The reaction typically is performed at temperatures between -10°C and room tempera-  
25 ture. A nonlimiting example of a solvent is dichloromethane. Nonlimiting examples of bases are triethylamine or di-isopropyl ethyl amine.

Scheme 5




Scheme 5 illustrates an alternative synthesis for compounds of formula 5. Compounds of formula 3 can be converted to compounds of formula 10 following procedures described in Scheme 2. Compounds of formula 10 then can be converted to compounds of formula 5 following procedures described in Scheme 4.


Scheme 6




11




12



13



4



As shown in Scheme 6, compounds of formula 11 can be converted to compounds of formula 12 by treatment with a base, such as potassium carbonate, triethylamine, or sodium hydride, followed by the addition of  $R^{11}X$ , wherein X is a halide, mesylate, or tosylate. Examples of solvents used in this reaction include DMF, THF,  $CH_2Cl_2$ , and mixtures thereof. The reaction is conducted at temperatures between  $0^\circ C$  and  $100^\circ C$  for about 15 minutes to 12 hours.

Alternatively, compounds of formula 11 can be admixed with a compound of formula  $R^{11}X$ , wherein X is hydroxyl, and the resulting mixture is treated with triphenylphosphine and diisopropylazodicarboxylate in a solvent, such as THF, to provide compounds of formula 12.


Compounds of formula 12 can be treated with hydrogen gas in the presence of a catalyst such as platinum oxide, palladium on carbon, or Raney nickel, or treated with an acid source, such as saturated aqueous ammonium chloride or aqueous hydrogen chloride in the presence of zinc metal, to provide compounds of formula 13. Examples of solvents used in this reaction include methanol, ethanol, ethyl acetate, or mixtures thereof. The reaction generally is conducted at room temperature or below for periods of one to twelve hours.

Compounds of formula 15 can be prepared by combining compounds of formula 13 with compounds of formula 4 (prepared as described in Scheme 1). Examples

- 44 -

of solvents used in this reaction include toluene, benzene, and xylene. The reaction is performed at temperatures of 60°C to 100°C for five to twelve hours.


Scheme 7



Scheme 7 shows an alternative synthesis of compounds of formula 15. Compounds of formula 3 are treated with compounds of formula 7, which is prepared according to Scheme 3. One solvent that can be used is DMF, and the reaction temperature is maintained between room temperature and 60°C over a one- to twelve-hour time period.


10

Scheme 8



5                    Scheme 8 shows an alternative approach to com-  
 pounds of formula 15. Compounds of formula 19 are con-  
 verted to compounds of formula 12 by treatment with an  
 alcohol in the presence of a base, such as sodium hy-  
 10    dride, potassium bis(trimethylsilyl)amide, or n-butyl-  
 lithium. Examples of solvents used in this reaction  
 include THF or diethyl ether. The reaction typically is  
 performed at temperatures between  $-15^\circ\text{C}$  and room tempera-  
 ture for about 1 to 6 hours. Compounds of formula 12 are

converted to compounds of formula 15 following procedures described in Scheme 6.



5                    Scheme 9 shows an alternative synthesis of  
 compounds 21 and 23. Compound 20 can be synthesized  
 using procedures described in Scheme 5 or Scheme 6,  
 wherein R<sub>2</sub>=Br. Compound 20 can be converted to Compound  
 21 by treatment with heteroaryl boronic acids (HetB(OH)<sub>2</sub>)  
 10 and a source of palladium(0) (for example, palladium  
 tetrakis triphenylphosphine) in the presence of a basic  
 aqueous solution, such as sodium bicarbonate, potassium  
 carbonate, or potassium phosphate. Nonlimiting examples  
 of solvents used in this reaction include THF, dioxane,  
 15 or ethylene glycol dimethyl ether. The reaction typ-  
 ically is performed at temperatures between 0°C and 90°C  
 for about 1 to 12 hours. Alternatively, an heteroaryl  
 boronic acid can be replaced with a heteroaryl stannate,  
 e.g., (HetSn(Bu)<sub>3</sub>).

20                    Compound 21 can also be converted to 22 by  
 treatment, for example, with zinc cyanide and a source of

- 47 -

palladium(0) (for example, palladium tetrakis triphenylphosphine), in the presence of a base such as triethylamine or Hunig's base. Nonlimiting examples of solvents used in this reaction are DMF and DME, at 80°C for 1 to 12 hours. Alternatively, compound 22 can be obtained from 21, using potassium cyanide, in the presence of a source of Pd(0) (such as palladium tetrakis triphenylphosphine), and a source of copper (such as copper iodide). The reaction is typically performed at temperatures between room temperature and 200°C, for about 30 minutes to 5 hours. Nonlimiting examples of solvents used in this reaction include DMF.

Finally, Compound 22 can be converted to Compound 23, using for example sodium azide, in the presence of a base such as triethylamine. Nonlimiting examples of solvents used in this reaction include DMF or nitrobenzene. The reaction typically is conducted at temperatures between 80°C and 100°C, for about 1 to 5 hours.

Specific, nonlimiting examples of compounds of structural formula (I) are provided below, the synthesis of which were performed in accordance with the procedures set forth below and in copending U.S. Patent Application Publication No. 2003-0069284 A1, incorporated herein by reference.


Abbreviations used in the syntheses described herein are: hours (h), water (H<sub>2</sub>O), magnesium sulfate (MgSO<sub>4</sub>), hydrochloric acid (HCl), dimethyl sulfoxide (DMSO), diisopropyl azodicarboxylate (DIAD), methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>), chloroform (CHCl<sub>3</sub>), methanol (MeOH), ammonium hydroxide (NH<sub>4</sub>OH), deuterated chloroform (CDCl<sub>3</sub>), tetrahydrofuran (THF), N-methylpyrrolidone (NMP), acetic acid (AcOH), sodium hydroxide (NaOH), ethyl acetate (EtOAc), ethanol (EtOH), dimethyl sulfoxide (DMSO), diethyl ether (Et<sub>2</sub>O), sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>), sodium



- 48 -

bicarbonate ( $\text{NaHCO}_3$ ), nitric acid ( $\text{HNO}_3$ ), sodium chloride ( $\text{NaCl}$ ), sodium sulfate ( $\text{Na}_2\text{SO}_4$ ), dimethylformamide (DMF), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), and N,N-diisopropylethylamine (DIEA).


5 Intermediate 1:



5-Methyl-pyrazine-2-carbonyl azide

To a stirred suspension of 5-methyl-pyrazine-2-carboxylic acid (25 g, 181 mmol) in 540 mL THF at room temperature under nitrogen was added DIEA (31.7 mL, 181 mmol) resulting in a brown solution. Diphenyl phosphoryl azide (39.2 mL, 181 mmol) then was added dropwise as a solution in 50 mL THF over 1 hour behind a blast shield. The reaction was allowed to stir overnight. The reaction then was rotary evaporated to a small volume at room temperature and partitioned between  $\text{Et}_2\text{O}$  (1 L) and  $\text{H}_2\text{O}$  (1 L). The  $\text{H}_2\text{O}$  layer was back extracted with 2 x 250 mL  $\text{Et}_2\text{O}$ , and the combined organics washed 2 x 1L with saturated sodium bicarbonate. The organics were dried ( $\text{MgSO}_4$ ), filtered, and concentrated to a solid mass, which was triturated with  $\text{Et}_2\text{O}$  to give the product as a yellow solid (15 g, 50%). Purer compound could be isolated by taking x g of the crude product in 20x mL of  $\text{Et}_2\text{O}$ , and treating with 1-2x g of decolorizing carbon at room temperature for a few minutes. After filtration and concentration, this material was homogeneous by TLC in  $\text{EtOAc}$  and pure white. The recovery was typically 65%.

- 49 -

Compound 1:

1-[2-(Piperidin-3-ylmethoxy)-5-trifluoromethyl-phenyl]-3-(5-trifluoromethyl-pyrazin-2-yl)-urea hydrochloride salt

5                    Step 1: 3-[2-(4-Nitro-phenoxy-carbonylamino)-4-trifluoromethyl-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester. To a stirred solution of 3-(2-amino-4-trifluoromethyl-phenoxy-methyl)-piperidine-1-carboxylic acid tert-butyl ester (240 mg, 0.64 mmol) in 2 mL CH<sub>2</sub>Cl<sub>2</sub>, at 0°C under nitrogen was added pyridine (57 uL, 0.71 mmol) followed by p-nitrophenyl chloroformate (130 mg, 0.64 mmol). After 1 hour at 0°C, the reaction mixture was diluted to 30 mL with CH<sub>2</sub>Cl<sub>2</sub>, then washed 2 x 30 mL with 2N HCl, 1 x 30 mL with water, and 1 x 30 mL with brine. The organics were dried (MgSO<sub>4</sub>), filtered, and concentrated to an off white foam.


20                    Step 2: 3-[4-Trifluoromethyl-2-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester. 3-[2-(4-Nitro-phenoxy-carbonylamino)-4-trifluoromethyl-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester (217 mg, 0.4 mmol) and 5-trifluoromethyl-pyrazin-2-ylamine (66 mg, 0.4 mmol) (prepared according to the method of U.S. 4,293,552) were mixed as solids in a 5 mL reaction vial, 25 diluted with 400 uL NMP, capped and stirred as a dark

- 50 -

yellow solution that then was immersed in an oil bath at 85°C and stirred for six hours. The reaction mixture was cooled to room temperature and stirred overnight. NMP then was removed by Kugelrohr distillation at 0.5 mm and 80°C. The brown residue was diluted to 30 mL with CH<sub>2</sub>Cl<sub>2</sub>, then washed 3 x 30 mL with 1M Na<sub>2</sub>CO<sub>3</sub> to remove p-nitro-phenol. The organics were dried (MgSO<sub>4</sub>), filtered, and concentrated to a crude solid that was triturated with EtOAc and filtered to give the desired product as a white solid.

Step 3: To a stirred solution of 3-[2-(4-nitro-phenoxy-carbonylamino)-4-trifluoromethyl-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester (60 mg, 0.11 mmol) in 2 mL dioxane at room temperature in a capped flask was added 2N HCl in dioxane (2 mL) and the reaction mixture was stirred overnight. The resulting suspension was concentrated by rotary evaporation and high vacuum to give a yellow solid corresponding to the final product (57 mg, 99%). <sup>1</sup>H-NMR (400 MHz, d<sub>6</sub>-DMSO) δ: 11.18 (br s, 1H), 9.91 (s, 1H), 9.13 (s, 1H), 8.79 (s, 1H), 8.62 (br s, 1H), 8.57 (s, 1H), 7.43 (d, 1H), 7.25 (d, 1H), 4.18 (m, 2H), 3.44 (m, 1H), 3.23 (m, 1H), 2.86 (m, 2H), 2.36 (m, 1H), 1.96 (m, 1H), 1.83 (m, 1H), 1.67 (m, 1H), 1.42 (m, 1H). LRMS (apci, positive) m/e 464.3 (M+1).


- 51 -

Compound 2:

1-[5-Methyl-2-(piperidin-3-ylmethoxy)-phenyl]-3-(5-trifluoromethyl-pyrazin-2-yl)-urea hydrochloride salt

5                    Step 1: 3-{4-Methyl-2-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-phenoxy-methyl}-piperidine-1-carboxylic acid tert-butyl ester. Prepared from 3-[4-methyl-2-(4-nitro-phenoxy-carbonylamino)-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester (WO  
10 02/070494) according to the procedure of Compound 1, Step 2. The product was isolated as a white solid.


                  Step 2: Prepared from 3-{4-methyl-2-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-phenoxy-methyl}-  
15 piperidine-1-carboxylic acid tert-butyl ester according to the procedure of Compound 1, Step 3. The final product was isolated as a yellow solid (30 mg, 99%). <sup>1</sup>H-NMR (400 MHz, d<sub>6</sub>-DMSO) δ: 10.86 (s, 1H), 9.61 (s, 1H), 9.05 (s, 1H), 8.77 (s, 1H), 8.58 (br s, 1H), 7.98 (s,  
20 1H), 6.98 (d, 1H), 6.83 (d, 1H), 3.99 (m, 2H), 3.40 (m, 1H), 3.23 (m, 1H), 2.81 (m, 2H), 2.23 (m, 1H), 2.22 (s, 3H), 1.91 (m, 1H), 1.80 (m, 1H), 1.62 (m, 1H), 1.39 (m, 1H). LRMS (apci, positive) m/e 410.4 (M+1).




Compound 3:


1-[5-Methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-trifluoromethyl-pyrazin-2-yl)-urea

5                   Prepared according to the procedure of  
Compound 1 Step 2 from [5-methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-carbamic acid 4-nitro-phenyl ester (WO 02/070494) as a white solid. <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ:  
11.01 (br s, 1H), 8.87 (br s, 1H), 8.82 (s, 1H), 8.44 (s,  
10 1H), 8.17 (s, 1H), 6.84 (d, 1H), 6.76 (d, 1H), 3.84 (m,  
2H), 3.16 (m, 1H), 2.81 (m, 1H), 2.37 (s, 3H), 2.29 (m,  
1H), 2.20 (s, 3H), 1.99 (m, 1H), 1.87-1.62 (m, 4H), 1.11  
(m, 1H). LRMS (apci, positive) m/e 424.3 (M+1).

Additional nonlimiting compounds of the  
15 present invention are illustrated below.

Compound 4

Compound 5Compound 65 Compound 7

Compound 8Therapeutic Methods

Compounds of the present invention can be used  
5 to potentiate the therapeutic effects of radiation and/or  
a chemotherapeutic agent used in the treatment of cancers  
and other cell proliferation indications involving  
eukaryotic cells, including those in humans and other  
animals. For example, compounds of the invention can be  
10 used to enhance treatment of tumors that are customarily  
treated with an antimetabolite, e.g., methotrexate or 5-  
fluorouracil (5-FU). In general, the present compounds  
inhibit aberrantly proliferating cells, both cancerous  
and noncancerous.

15 Use of compounds of the present invention can  
result in partial or complete regression of aberrantly  
proliferating cells, i.e., the partial or complete  
disappearance of such cells from the cell population.  
Thus, for example, when the population of aberrantly  
20 proliferating cells are tumor cells, the method of the  
invention can be used to slow the rate of tumor growth,  
decrease the size or number of tumors, or to induce  
partial or complete tumor regression.

In all embodiments, the invention can be used  
25 *in vivo* or *ex vivo* where no aberrant cell proliferation  
has been identified or where no aberrant cell prolifera-

tion is ongoing, but where aberrant cell proliferation is suspected or expected, respectively. Moreover, the invention also can be used wherever aberrant cell proliferation has been previously treated to prevent or inhibit recurrence of the same. In these and related embodiments, the "cell population comprising aberrantly proliferating cells" refers to any cell population where no aberrant cell proliferation has been identified or is ongoing, but where aberrant cell proliferation is suspected or expected, respectively, and/or any cell population previously treated for aberrant cell proliferation to prevent or inhibit recurrence of the same.

One method of the present invention comprises administration of a therapeutically effective amount of a present Chk1 inhibitor compound in combination with a chemotherapeutic agent that can effect single- or double-strand DNA breaks or that can block DNA replication or cell proliferation. Alternatively, a method of the present invention comprises administration of a therapeutically effective amount of at least one of the present Chk1 inhibitor compounds in combination with therapies that include use of an antibody, e.g., herceptin, that has activity in inhibiting the proliferation of cancer cells. Accordingly, cancers, for example, colorectal cancers, head and neck cancers, pancreatic cancers, breast cancers, gastric cancers, bladder cancers, vulvar cancers, leukemias, lymphomas, melanomas, renal cell carcinomas, ovarian cancers, brain tumors, osteosarcomas, and lung carcinomas, are susceptible to enhanced treatment by administration of a present Chk1 inhibitor in combination with a chemotherapeutic agent or an antibody.

Cancers include tumors or neoplasms which are growths of tissue cells wherein multiplication of cells is uncontrolled and progressive. Some such growths are



- 56 -

benign, but others are termed "malignant," and can lead to death of the organism. Malignant neoplasms, or "cancers," are distinguished from benign growths in that, in addition to exhibiting aggressive cellular proliferation, can invade surrounding tissues and metastasize. Moreover, malignant neoplasms are characterized by showing a greater loss of differentiation (greater "dedifferentiation") and organization relative to one another and surrounding tissues. This property is called "anaplasia."

10           Cancers treatable by the present invention also include solid tumors, i.e., carcinomas and sarcomas. Carcinomas include malignant neoplasms derived from epithelial cells which infiltrate (i.e., invade) surrounding tissues and give rise to metastases. Adenocarcinomas are  
15           carcinomas derived from glandular tissue, or from tissues that form recognizable glandular structures. Another broad category of cancers includes sarcomas, which are tumors whose cells are embedded in a fibrillar or homogeneous substance, like embryonic connective tissue. The  
20           present invention also enables treatment of cancers of the myeloid or lymphoid systems, including leukemias, lymphomas, and other cancers that typically are not present as a tumor mass, but are distributed in the vascular or lymphoreticular systems.

25           Chk1 activity is associated with various forms of cancer in, for example, adult and pediatric oncology, growth of solid tumors/malignancies, myxoid and round cell carcinoma, locally advanced tumors, metastatic cancer, human soft tissue sarcomas, including Ewing's  
30           sarcoma, cancer metastases, including lymphatic metastases, squamous cell carcinoma, particularly of the head and neck, esophageal squamous cell carcinoma, oral carcinoma, blood cell malignancies, including multiple myeloma, leukemias, including acute lymphocytic leukemia,

- 57 -

acute nonlymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, and hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer (including small cell carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, 5 non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, nonsmall cell cancers, breast cancer, including small cell carcinoma and ductal carcinoma), gastrointestinal cancers (including stomach 10 cancer, colon cancer, colorectal cancer, and polyps associated with colorectal neoplasia), pancreatic cancer, liver cancer, urological cancers (including bladder cancer, such as primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, and 15 muscle-invasive bladder cancer), prostate cancer, malignancies of the female genital tract (including ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid 20 tumors in the ovarian follicle), malignancies of the male genital tract (including testicular cancer and penile cancer), kidney cancer (including renal cell carcinoma, brain cancer (including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, and 25 metastatic tumor cell invasion in the central nervous system), bone cancers (including osteomas and osteosarcomas), skin cancers (including malignant melanoma, tumor progression of human skin keratinocytes, and squamous cell cancer), thyroid cancer, retinoblastoma, neuro- 30 blastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma. Accordingly, administration of a present Chk1 inhibitor is expected to enhance treatment regimens.

- 58 -

Compounds of the present invention also can potentiate the efficacy of drugs in the treatment of inflammatory diseases. Examples of diseases that can benefit from combination therapy with compounds suitable for the method of the present invention are rheumatoid arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus erythematosus (SLE). Treatment of arthritis, Wegener's granulomatosis, and SLE often involves the use of immunosuppressive therapies, such as ionizing radiation, methotrexate, and cyclophosphamide. Such treatments typically induce, either directly or indirectly, DNA damage. Inhibition of Chk1 activity within the offending immune cells render the cells more sensitive to control by these standard treatments.

Psoriasis and vitiligo commonly are treated with ultraviolet radiation (UV) in combination with psoralen. The present DNA damaging agents induce the killing effect of UV and psoralen, and increase the therapeutic index of this treatment regimen. In general, compounds useful in methods of the present invention potentiate control of inflammatory disease cells when in combination with currently used immunosuppressive drugs.

In addition to the cancers disclosed above, the present invention also can be used in methods of treating noncancerous proliferating cells. Such conditions include, but are not limited to, atherosclerosis, restenosis, vasculitis, nephritis, retinopathy, renal disease, proliferative skin disorders, psoriasis, keloid scarring, actinic keratosis, Stevens-Johnson Syndrome, rheumatoid arthritis (RA), systemic-onset juvenile chronic arthritis (JCA), osteoporosis, systemic lupus erythematosus, hyperproliferative diseases of the eye including epithelial down growth, proliferative vitreo-

- 59 -

retinopathy (PVR), diabetic retropathy, Hemangio-proliferative diseases, ichthyosis, or papillomas.

One preferred method of administering a Chk1 inhibitor of the present invention is described in Keegan  
5 et al., U.S. Provisional application no. 60/503,925, filed September 17, 2003, the entire disclosure of which is incorporated herein by reference herein. Such methods for inhibiting aberrant cell proliferation involve scheduling the administration of a Chk1 activator (e.g.,  
10 a chemotherapeutic agent) and a Chk1 inhibitor according to the present invention. In this method, at least one Chk1 activator is administered at a dose and for a time sufficient to induce substantial synchronization of cell cycle arrest in proliferating cells. Upon achieving  
15 substantial phase synchronization, at least one Chk1 inhibitor is administered to abrogate the cell cycle arrest and induce therapeutic cell death. The method is useful with any Chk1 activator, and finds application in treating or preventing cancerous and noncancerous aberrant  
20 cell proliferation.

A population of aberrantly proliferating cells can be contacted with one Chk1 inhibitor or can be contacted with more than one Chk1 inhibitor. If more than one Chk1 inhibitor is used, the Chk1 inhibitors can be  
25 coadministered or administered at separate times as determined by the attending physician or laboratory technician.

A population of aberrantly proliferating cells also can be contacted with one Chk1 activator or can be  
30 contacted with more than one Chk1 activator. If more than one Chk1 activator is used, the Chk1 activators can be coadministered or administered at separate times as determined by the attending physician or laboratory technician.

- 60 -

The Chk1 inhibitors of the present invention can be employed to study or modify the behavior of cell populations *ex vivo*. For example, the present compounds can be used *ex vivo* to determine the optimal schedule.

5 and/or dosing of administration of a Chk1 inhibitor for a given indication, cell type, patient, and other parameter. Information gleaned from such use can be used for experimental purposes or in the clinic to set protocol for *in vitro* treatment. Other *ex vivo* uses for which the  
10 invention is suited are apparent to those skilled in the art.

A Chk1 inhibitor compound of the present invention also can radiosensitize a cell. Diseases that are treatable with electromagnetic radiation include  
15 neoplastic diseases, benign and malignant tumors, and cancerous cells.

Electromagnetic radiation treatment of other diseases not listed herein also is contemplated by the present invention. Preferred embodiments of the present  
20 invention employ the electromagnetic radiation of: gamma radiation ( $10^{-20}$  to  $10^{-13}$  m), X-ray radiation ( $10^{-12}$  to  $10^{-19}$  m), ultraviolet light (10 nm to 400 nm), visible light (400 nm to 700 nm), infrared radiation (700 nm to 1.0 mm), and microwave radiation (1 mm to 30 cm).

25 Many cancer treatment protocols currently employ radiosensitizers activated by electromagnetic radiation, e.g., X-rays. Examples of X-ray-activated radiosensitizers include, but are not limited to, the following: metronidazole, misonidazole, desmethyl-  
30 misonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, EO9, RB 6145, nicotinamide, 5-bromodeoxyuridine (BUdR), 5-iododeoxyuridine (IUdR), bromodeoxycytidine, fluorodeoxyuridine (FUdR),

- 61 -

hydroxyurea, cis-platin, and therapeutically effective analogs and derivatives of the same.

Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, PHOTOFRIN<sup>®</sup>, benzoporphyrin derivatives, NPe6, tin etioporphyrin (SnET2), pheoborbide-a, bacteriochlorophyll-a, naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

Radiosensitizers can be administered in conjunction with a therapeutically effective amount of one or more compounds in addition to the Chk1 inhibitor, such compounds including, but not limited to, compounds that promote the incorporation of radiosensitizers to the target cells, compounds that control the flow of therapeutics, nutrients, and/or oxygen to the target cells, chemotherapeutic agents that act on the tumor with or without additional radiation, or other therapeutically effective compounds for treating cancer or other disease. Examples of additional therapeutic agents that can be used in conjunction with radiosensitizers include, but are not limited to, 5-fluorouracil (5-FU), leucovorin, oxygen, carbogen, red cell transfusions, perfluorocarbons (e.g., FLUOSOL<sup>®</sup>-DA), 2,3-DPG, BW12C, calcium channel blockers, pentoxifylline, antiangiogenesis compounds, hydralazine, and L-BSO.

Chemotherapeutic agents that can be used include, but are not limited to, alkylating agents, anti-metabolites, hormones and antagonists thereof, radioisotopes, antibodies, as well as natural products, and combinations thereof. For example, an inhibitor compound of the present invention can be administered with antibi-

- 62 -

otics, such as doxorubicin and other anthracycline analogs, nitrogen mustards, such as cyclophosphamide, pyrimidine analogs such as 5-fluorouracil, cis-platin, hydroxyurea, taxol and its natural and synthetic derivatives, and the like. As another example, in the case of mixed tumors, such as adenocarcinoma of the breast, where the tumors include gonadotropin-dependent and gonadotropin-independent cells, the compound can be administered in conjunction with leuprolide or goserelin (synthetic peptide analogs of LH-RH). Other antineoplastic protocols include the use of an inhibitor compound with another treatment modality, e.g., surgery or radiation, also referred to herein as "adjunct anti-neoplastic modalities." Additional chemotherapeutic agents useful in the invention include hormones and antagonists thereof, radioisotopes, antibodies, natural products, and combinations thereof. Examples of chemotherapeutic agents useful for the method of the present invention are listed in the following table.

Alkylating agents	Nitrogen mustards	mechlorethamine
cyclophosphamide	ifosfamide	melphalan
chlorambucil	Nitrosoureas	carmustine (BCNU)
lomustine (CCNU)	semustine (methyl-CCNU)	Ethylenimine/Methylmelamine
thiethylenemelamine (TEM)	triethylene thiophosphoramide	(thiotepa)
hexamethylmelamine	(HMM, altretamine)	Alkyl sulfonates
busulfan	Triazines	dacarbazine (DTIC)
Antimetabolites	Folic Acid analogs	methotrexate
trimetrexate	Pyrimidine analogs	5-fluorouracil
fluorodeoxyuridine	gemcitabine	cytosine arabinoside
(AraC, cytarabine)	5-azacytidine	2,2'-difluorodeoxycytidine
Purine analogs	6-mercaptopurine	6-thioguanine
azathioprine	2'-deoxycoformycin	(pentostatin)
erythrohydroxynonyladenine (EHNA)	fludarabine phosphate	2-chlorodeoxyadenosine
(cladribine, 2-CdA)	multitargeted antifolate	Type I Topoisomerase Inhibitors
camptothecin	topotecan	irinotecan
Natural products	Antimitotic drugs	paclitaxel
Vinca alkaloids	vinblastine (VLB)	vincristine
vinorelbine	Taxotere (docetaxel)	estramustine
estramustine phosphate	etoposide	teniposide
Epipodophylotoxins		
Antibiotics	actinomycin D	daunomycin (rubidomycin)
doxorubicin (adriamycin)	mitoxantroneidarubicin	bleomycinsplicamycin (mithramycin)
mitomycinC	dactinomycin	Enzymes
L-asparaginase	Biological response modifiers	interferon-alpha
IL-2	G-CSF	GM-CSF
Differentiation Agents	retinoic acid derivatives	Radiosensitizers
metronidazole	misonidazole	desmethylmisonidazole
pimonidazole	etanidazole	nimorazole
RSU 1069	EO9	RB 6145
SR4233	nicotinamide	5-bromodeoxyuridine
5-iododeoxyuridine	bromodeoxycytidine	Miscellaneous agents
Platinum coordination complexes	cis-platin	carboplatin
oxaliplatin	Anthracenedione	mitoxantrone
Substituted urea	hydroxyurea	Methylhydrazine derivatives
N-methylhydrazine (MIH)	procabazine	Adrenocortical suppressant
mitotane (o,p'-DDD)	ainoglutethimide	Cytokines
interferon ( $\alpha$ , $\beta$ , $\gamma$ )	interleukin-2	Adrenocorticosteroids/antagonists
	Hormones and antagonists	
prednisone and equivalents	dexamethasone	ainoglutethimide
Progestins	hydroxyprogesterone caproate	medroxyprogesterone acetate
megestrol acetate	Estrogens	diethylstilbestrol
ethynyl estradiol/ equivalents	Antiestrogen	tamoxifen
Androgens	testosterone propionate	fluoxymesterone/equivalents
Antiandrogens	flutamide	gonadotropin-releasing
hormone analogs	leuprolide	Nonsteroidal antiandrogens
flutamide	Photosensitizers	hematoporphyrin derivatives
Photofrin	benzoporphyrin derivatives	Npe6
tin etioporphyrin (SnET2)	pheoboride-a	bacteriochlorophyll-a
naphthalocyanines	phthalocyanines	zinc phthalocyanines
Growth Factor Receptor Antagonists	EGFR antagonists	HER-2 antagonists



- 64 -

Examples of chemotherapeutic agents that are particularly useful in conjunction with radiosensitizers include, for example, camptothecin, carboplatin, cis-platin, daunorubicin, doxorubicin, interferon (alpha, 5 beta, gamma), irinotecan, hydroxyurea, chlorambucil, 5-fluorouracil (5-FU), methotrexate, 2-chloroadenosine, fludarabine, azacytidine, gemcitabine, pemetrexed, interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, and therapeutically effective analogs and derivatives of 10 the same.

In accordance with the present invention, compounds of the present invention are useful in combination with gemcitabine, or in combination with gemcitabine and paclitaxel. Compounds of the present invention also are 15 useful in combination with pemetrexed, or in combination with pemetrexed and cisplatin, carboplatin, or other platins. A present Chk1 inhibitor also can be administered in combination with gemcitabine and pemetrexed.

A present Chk1 inhibitor administered in combination with gemcitabine can be useful in the treatment 20 of, for example, pancreatic carcinoma, leiomyosarcoma of the uterus, bone sarcoma, metastatic non-small cell lung cancer, extremity and trunk soft tissue sarcoma, renal cell cancer, adenocarcinoma, and Hodgkin's disease. A 25 present Chk1 inhibitor administered with pemetrexed can be useful in the treatment of mesothelioma.

As appreciated by persons skilled in the art, reference herein to treatment extends to prophylaxis, as well as to treatment of established diseases or symptoms. 30 Reference to treatment also refers to the reduction of the rate of proliferation or the reduction of recurrence of the treated indication. It is further appreciated that the amount of a compound of the invention required for use in treatment varies with the nature of the condi-

- 65 -

tion being treated, and with the age and the condition of the patient, and is ultimately determined by the attendant physician or veterinarian.

In general, however, doses administered for adult human treatment typically are in the range of 0.001 mg/kg to about 100 mg/kg per day. The desired dose can be conveniently administered in a single dose, or as multiple doses administered at appropriate intervals, for example as two, three, four or more subdoses per day. In practice, the physician determines the actual dosing regimen most suitable for an individual patient, and the dosage varies with the age, weight, and response of the particular patient. The above dosages are exemplary of the average case, but individual instances can exist wherein higher or lower dosages are merited, and such are within the scope of the present invention.

Contact of the cell population with a present Chk1 inhibitor can likewise occur at any dose and time sufficient to achieve substantial abrogation of the cell cycle checkpoint. Typically, though not necessarily, such times include up to about 72 to about 96 hours, depending upon various factors. In some embodiments, it is desirable or necessary to administer Chk1 inhibitor over a period of up to about several weeks or more, as determined by the attending physician or technician. Thus, a present Chk1 inhibitor typically can be administered for up to about 1 hour, up to about 2 hours, up to about 3 hours, up to about 4 hours, up to about 6 hours, up to about 12 hours, up to about 18 hours, up to about 24 hours, up to about 48 hours, or up to about 72 hours. Persons skilled in the art appreciate that the ranges of time expressed herein are merely exemplary and that ranges and subranges within and outside those expressed also are within the scope of the invention.

- 66 -

Chk1 inhibitors of the present invention can be administered over a plurality of doses. For example, the Chk1 inhibitor can be given at a frequency of: four doses delivered as one dose per day at four-day intervals (q4d x 4); four doses delivered as one dose per day at three-day intervals (q3d x 4); one dose delivered per day at five-day intervals (qd x 5); one dose per week for three weeks (qwk3); five daily doses, with two days rest, and another five daily doses (5/2/5); or, any dose regimen determined to be appropriate for the circumstance.

### EXAMPLES

#### Example 1

##### Determination of IC<sub>50</sub> Values of the Chk1 Inhibitors

Human Chk1 cDNA was identified and cloned as described previously in International Application Publication No. WO 99/11795, filed September 4, 1998. A FLAG<sup>®</sup> tag was inserted in frame with the amino terminus of the full-length Chk1. The 5' primer contains an EcoRI site, a Kozak sequence, and also encodes a FLAG<sup>®</sup> tag for affinity purification using the M2 Antibody (Sigma, Saint Louis, IL). The 3' primer contains a SalI site. The PCR-amplified fragment was cloned into pCI-Neo as an EcoRI-SalI fragment (Invitrogen, Carlsbad, CA), then subcloned as an EcoRI-NotI fragment into pFastBacI (Gibco-BRL, Bethesda, MD). Recombinant baculovirus was prepared as described in the Gibco-BRL Bac-to-Bac manual and used to infect Sf-9 cells grown in CCM3 medium (HyClone Laboratories, Logan, UT) for expression of FLAG<sup>®</sup>-tagged Chk1 protein.

FLAG<sup>®</sup>-tagged Chk1 was purified from frozen pellets of baculovirus-infected SF9 cells. Frozen cell

- 67 -

pellets were mixed with an equal volume of 2X lysis buffer containing 100 mM Tris-HCl pH 7.5, 200 mM NaCl, 50 mM B-glycerophosphate, 25 mM NaF, 4 mM MgCl<sub>2</sub>, 0.5 mM EGTA, 0.2% TWEEN<sup>®</sup>-20, 2 mM sodium vanadate, 2 mM DTT, and  
5 a cocktail of protease inhibitors (Complete mini, Boehringer Mannheim 2000 catalog #1836170). Cells then were dounced 20 times with the loose pestle of a dounce homogenizer and centrifuged at 48,400 x g for 1 hour. The M2 affinity was prewashed with 10 column volumes of  
10 50 mM glycine pH 3.5 followed by 20 mM Tris pH 7.5, 150 mM NaCl alternating three times and ending with a Tris NaCl wash. The column then was washed with 25 column volumes of 20 mM Tris pH 7.5, 150 mM NaCl, 0.1% TWEEN<sup>®</sup>-20, 1 mM EGTA, 1 mM EDTA and 1X complete mini protease  
15 tablets. The cleared lysate then was bound to M2 affinity resin in batch at 4°C for 4 hours. The mixture of resin and lysate then was poured into a column and the flow through collected. The resin was washed with 10 column volumes of 20 mM Tris pH 7.5, 150 mM NaCl, and 3  
20 mM N-octyl glucoside. FLAG<sup>®</sup>-tagged Chk1 then was eluted from the column with 6 column volumes of cold 20 mM Tris pH 7.5, 150 mM NaCl, 3 mM N-octyl glucoside containing 0.5 mg/mL FLAG<sup>®</sup> peptide (Sigma, 2000 Catalog # F-3290). Three fractions were collected and analyzed for the pres-  
25 ence of FLAG-tagged Chk1.

The protein kinase was used in an assay for Chk1 kinase activity that includes 100 ng purified FLAG<sup>®</sup>-Chk1 (150 pmol of ATP/min), 20 μM Cdc25C peptide (H-leu-tyr-arg-ser-pro-ser-met-pro-glu-asn-leu-asn-arg-arg-arg-  
30 arg-OH) (SEQ ID NO: 1), 4 μM ATP, 2 μCi [<sup>32</sup>P]γ-ATP, 20 mM Hepes pH 7.2, 5 mM MgCl<sub>2</sub>, 0.1% NP40, and 1 mM DTT. This assay was used to determine IC<sub>50</sub> of compounds of the present invention. Reactions were initiated by the addition of ATP-containing reaction mix and carried out at room

- 68 -

temperature for 10 min. Reactions were stopped by the addition of phosphoric acid (150 mM final concentration) and transferred to phosphocellulose discs. The phosphocellulose discs were washed five times with 150 mM phosphoric acid and air-dried. Scintillation fluid was added and discs were counted in a Wallac scintillation counter. Chk1 inhibitors of the present invention that were subjected to the assay have measured IC<sub>50</sub> values of about 8 to about 500 nM.

## 10 Example 2

### Selectivity

Chk1 inhibitors of the present invention were tested for selectivity as against one or more other protein kinases, i.e., DNA-PK, Cdc2, Casein Kinase I (CKI), Chk2, p38 MAP kinase, ERK kinase, Protein Kinase A (PKA), and/or calcium-calmodulin protein kinase II (CaM KII). Assay procedures for all of these kinases except Chk2 have been previously described in the literature, including U.S. Patent Publication No. 2002-016521 A1, and U.S. patent application 08/184,605, filed January 21, 1994, both of which are herein incorporated by reference.

Activity of the compounds against Chk2 was assayed as follows: 128 ng of purified His-tagged Chk2 was incubated with up to 100 mM Chk1 inhibitor in the presence of 4 mM ATP, 1 mCi [<sup>32</sup>P]γ-ATP, 20 mM Hepes pH 7.5, 5 mM MgCl<sub>2</sub>, and 0.25% NP40 for 20 minutes at room temperature. Reactions were stopped with a final concentration of 150 mM phosphoric acid, and 5/8 of the reaction mixture was transferred to phosphocellulose discs. The discs were washed five times with 150 mM phosphoric acid, and air-dried. Scintillant was added

- 69 -

and radioactivity was counted using a Wallac beta counter.

p38 MAP kinase, ERK kinase, PKA, CaM KII, and Cdc2 were purchased from New England Biolabs, and assays were performed according to the manufacturer's instructions using 4-50  $\mu$ M ATP and testing Chk1 inhibitor concentrations as high as 100  $\mu$ M. All inhibitors tested showed at least a 100-fold selectivity for Chk1 over the other enzymes.

### 10 Example 3

#### Chk1 Inhibitors of the Invention Inhibit Chk1 Function in Cells

To establish that the Chk1 inhibitors of the invention inhibit Chk1 function in cells, inhibitors can be tested in molecular cell-based assays. Because mammalian Chk1 has been shown to phosphorylate Cdc25C in vitro, suggesting that it negatively regulates cyclin B/cdc2 in response to DNA damage, the ability of the Chk1 inhibitors to enhance the activity of CyclinB/cdc2 can be analyzed. The experiment can be designed as follows: HeLa cells are irradiated with 800 rads and incubated for 7 hours at 37°C. Because these cells are functionally p53 negative, they arrest exclusively in G2. Then, nocodazole is added to a concentration of 0.5  $\mu$ g/mL and the cells are incubated for 15 hours at 37°C. The addition of nocodazole is designed to trap any cells that progress through the G2 arrest into M. Finally, a Chk1 inhibitor is added for 8 hours, the cells harvested, lysed and immunoprecipitated equal amounts of protein with an antibody to Cyclin B1 (New England Biolabs) as suggested by the manufacturer. Immunoprecipitates then are analyzed for Cyclin B-associated cdc2 kinase activity

- 70 -

by assaying histone H1 kinase activity (Yu et al., *J Biol Chem.*, Dec. 11, 1998; 273(50):33455-64).

In addition, the ability of the subject Chk1 inhibitors to abrogate the ionizing radiation-induced G2 DNA damage checkpoint can be established using mitotic index assay experiments. HeLa cells (approximately  $1 \times 10^6$ ) are treated as described above. Cells are harvested by centrifugation, washed once with PBS, then resuspended in 2.5 mL of 75 mM KCl and centrifuged again. The cells then are fixed in 3 mL of freshly prepared cold acetic acid:methanol (1:3) and incubated on ice for 20 minutes. Cells are pelleted, fix solution aspirated and resuspended in 0.5 mL of PBS. Mitotic spreads are prepared by pipeting 100  $\mu$ L of the fixed cells onto a glass microscope slide and flooding the sample with 1 mL of fix solution. Slides then are air dried, stained with Wright's stain (Sigma) for 1 minute, followed by one wash with water and one wash with 50% methanol. The presence of condensed chromosomes and lack of nuclear envelope identifies mitotic cells.

#### Example 4

##### Chk1 Inhibitors of the Present Invention Enhance Killing of Cells by Cancer Treatments

To demonstrate that the inhibition of Chk1 by a compound of the present invention sensitizes targeted cells to the killing effect of DNA-damaging agents, cells can be incubated in the presence of a present Chk1 inhibitor and exposed to either irradiation or a chemical DNA-damaging agent. Cells plated at a density of 1000-2000 per well in 96-well microtitre plates are grown in RMPI 1640 containing 10% FBS, 100 U/mL penicillin and 100  $\mu$ g/mL streptomycin for 18 hours at 37°C in a humidified

- 71 -

- incubator with 5% CO<sub>2</sub>. Cells tested can include any cells or cell lines of interest, such as HeLa, ACHN, 786-0, HCT116, SW620, HT29, Colo205, SK-MEL-5, SK-MEL-28, A549, H322, OVCAR-3, SK-OV-3, MDA-MB-231, MCF-7, PC-3, HL-60, K562, and MOLT4. All cell line designations refer to the following human cell lines:

HeLa	cervical adenocarcinoma
ACHN	renal adenocarcinoma
786-0	renal adenocarcinoma
HCT116	colon carcinoma
SW620	colon carcinoma, lymph node metastasis
HT-29	colonrectal adenocarcinoma
Colo205	colon adenocarcinoma
SK-MEL-5	melanoma
SK-MEL-28	malignant melanoma
A549	lung carcinoma
H322	broncholoalveolar carcinoma
OVCAR-3	ovarian adenocarcinoma
SK-OV-3	ovarian adenocarcinoma
MDA-MB-231	breast adenocarcinoma
MCF-7	breast adenocarcinoma
PC-3	prostate adenocarcinoma, from metastasis to bone
HL-60	acute promyelocytic leukemia
K562	chronic myelogenous leukemia
MOLT4	acute lymphoblastic leukemia; T lymphoblast

- Cells are treated with media containing chemotherapeutic drugs alone or chemotherapeutic drugs and a Chk1 inhibitor. Cells are incubated for approximately 5 days before growth is measured by determination of levels of <sup>3</sup>H-thymidine uptake. Chemotherapeutic drugs include etoposide, doxorubicin, cis-platin, chlorambucil, 5-fluorouracil (5-FU). The drug concentration necessary to inhibit cell growth to 90% of untreated control cells is defined as the GI<sub>90</sub>.



- 72 -

Compounds of the present invention can be tested with additional antimetabolites, including methotrexate, hydroxyurea, 2-chloroadenosine, fludarabine, azacytidine, and gemcitabine to assess therein ability to enhance killing of the agents. Compounds of the present invention can be compared to one another by assessing enhanced killing of HT29 colorectal carcinoma in combination with gemcitabine.

In addition, the ability of the Chk1 inhibitors of the invention to enhance killing by radiation can be tested.

#### Example 5

##### Animal Tumor Models

To test the ability of the Chk1 inhibitors of the invention to enhance the killing of tumors by DNA damaging agents in mice, xenograft tumor models using colon tumor cell lines are established. 5-fluorouracil (5-FU) or gemcitabine can be used as DNA damaging agents. HT29 and Colo205 (human colon carcinoma) and H460 and Calu-6 (nonsmall cell carcinoma) cells can be used to propagate xenograft tumors in 6-8 week old female thymic Balb/c (nu/nu) mice. Mice are maintained in a laminar airflow cabinet under pathogen-free conditions and fed sterile food and water *ad libitum*. Cell lines are grown to subconfluence in RPMI 1640 media supplemented with 10% FBS, 100 U/mL penicillin, 100  $\mu$ g/mL streptomycin, and 1.5 mM L-glutamine in a 5% CO<sub>2</sub> humidified environment. Single cell suspensions are prepared in CMF-PBS, and cell concentration adjusted to  $1 \times 10^8$  cells/mL. Mice are inoculated subcutaneously (s.c.) on the right flank or right leg with a total of  $1 \times 10^7$  cells (100  $\mu$ L).

- 73 -

Mice are randomized (5-15 mice/group) into four treatment groups and used when tumors reach a volume of 75-100 cm<sup>3</sup> (usually 7-11 days post-inoculation). Tumors are measured with vernier calipers and tumor


5 volumes are estimated using the empirically derived formula: tumor volume (cm<sup>3</sup>)=tumor length (cm) x tumor width (cm) x tumor depth (cm)/3.3. Treatment consists of i) 100 µL intraperitoneal (i.p) injection of gemcitabine at 160 mg/kg. A delay in tumor growth is observed in the

10 mice treated with gemcitabine. Treatment of mice with both 160 mg/kg gemcitabine in combination with oral administration of Chk1 inhibitors is expected to reduce tumor volumes and prolong life. Tumor size is monitored every other day for the duration of the experiment.

15 Obviously, many modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and, therefore, only such limitations should be imposed as are indicated by the appended claims.

WHAT IS CLAIMED IS:

1. A compound having a formula



wherein  $X^1$  is null, -O-, -S-, -CH<sub>2</sub>-, or -N(R<sup>1</sup>)-;

$X^2$  is -O-, -S-, or -N(R<sup>1</sup>)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C<sub>1-6</sub>alkyl substituted with a heteroaryl or aryl group, wherein (a) said aryl or heteroaryl group of group W is substituted with at least one of CF<sub>3</sub> and heteroaryl, (b) said aryl group of group W is optionally substituted with one to three substituents represented by R<sup>2</sup>, and (c) said heteroaryl group of group W is optionally substituted with one to three substituents represented by R<sup>5</sup>;

R<sup>1</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, and aryl;


R<sup>2</sup> is selected from the group consisting of heteroaryl, halo, optionally substituted C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, OCF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneOR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneSO<sub>2</sub>NR<sup>3</sup>, C<sub>1-6</sub>alkyleneOR<sup>3</sup>, and SR<sup>3</sup>;

- 75 -

$R^3$  is selected from the group consisting of hydro, halo,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl, cycloalkyl, aryl, heteroaryl,  $CO_2R^4$ ,  $SO_2R^4$ ,  $C_{1-6}$ alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl,  $N(R^4)_2$ , and  $SO_2R^4$ ,  $C_{1-6}$ alkylenearyl,  $C_{1-6}$ alkyleneheteroaryl,  $C_{1-6}$ alkylene $C_{3-8}$ heterocycloalkyl,  $C_{1-6}$ alkylene- $SO_2$ aryl, optionally substituted  $C_{1-6}$ alkylene $N(R^4)_2$ ,  $OCF_3$ ,  $C_{1-6}$ alkylene $N(R^4)_3^+$ ,  $C_{3-8}$ heterocycloalkyl, and  $CH(C_{1-6}alkyleneN(R^4)_2)_2$ , or two  $R^3$  groups are taken together to form an optionally substituted 3- to 6-membered aliphatic ring;

$R^4$  is selected from the group consisting of hydro,  $C_{1-6}$ alkyl, cycloalkyl, aryl, heteroaryl,  $C_{1-6}$ alkylenearyl, and  $SO_2C_{1-6}alkyl$ , or two  $R^4$  groups are taken together to form an optionally substituted 3- to 6-membered ring;

$R^5$  is selected from the group consisting of  $C_{1-6}$ alkyl, aryl, heteroaryl, heterocycloalkyl,  $N(R^3)_2$ ,  $OR^3$ , halo,  $N_3$ , CN,  $C_{1-6}alkylenearyl$ ,  $C_{1-6}alkyleneN(R^3)_2$ ,  $C(O)R^3$ ,  $C(O)OR^3$ ,  $C(O)N(R^3)_2$ ,  $N(R^1)C(O)R^3$ ,  $N(R^1)C(O)OR^3$ ,  $CF_3$ , and



$R^6$  is selected from the group consisting of  $OR^{11}$ ,  $-C\equiv C-R^7$ , and heteroaryl;

$R^7$  is selected from the group consisting of hydro,  $C_{1-6}$ alkyl, aryl,  $C_{1-6}alkylenearyl$ , heteroaryl,  $C_{1-6}alkyleneheteroaryl$ , and alkoxy;

$R^8$ ,  $R^9$ , and  $R^{10}$ , independently, are selected from the group consisting of hydro, halo, optionally

- 76 -

substituted C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, OCF<sub>3</sub>, CF<sub>3</sub>, NO<sub>2</sub>, CN, NC, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, CO<sub>2</sub>R<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, C(O)R<sup>3</sup>, N(R<sup>1</sup>)COR<sup>3</sup>, N(R<sup>1</sup>)C(O)OR<sup>3</sup>, N(R<sup>8</sup>)C(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkylene-C(O)R<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-3</sub>alkylene-OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkyleneNHC(O)OR<sup>3</sup>, N(R<sup>1</sup>)C(O)C<sub>1-6</sub>alkylene-SO<sub>2</sub>NR<sup>3</sup>, C<sub>1-6</sub>alkyleneOR<sup>3</sup>, and SR<sup>3</sup>;

R<sup>11</sup> is selected from the group consisting of hydro, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, N(R<sup>4</sup>)<sub>2</sub>, and SO<sub>2</sub>R<sup>4</sup>, C<sub>1-6</sub>alkylenearyl, C<sub>1-6</sub>alkyleneheteroaryl, C<sub>1-6</sub>alkyleneC<sub>3-8</sub>heterocycloalkyl, C<sub>1-6</sub>alkyleneSO<sub>2</sub>aryl, optionally substituted C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>2</sub>, OCF<sub>3</sub>, C<sub>1-6</sub>alkylene-N(R<sup>4</sup>)<sub>3</sub><sup>+</sup>, C<sub>3-8</sub>heterocycloalkyl, and CH(C<sub>1-6</sub>alkyleneN(R<sup>4</sup>)<sub>2</sub>)<sub>2</sub>;

and a pharmaceutically acceptable salt, or prodrug, or solvate thereof.

- 77 -

2. The compound of claim 1 wherein


X<sup>1</sup> and X<sup>2</sup> are -N(H)-;

Y is O or S;


W is heteroaryl containing at least two heteroatoms selected from the group consisting of N, O, and S, said ring optionally substituted with one or two substituents selected from the group consisting of C<sub>1-6</sub>alkyl, aryl, heteroaryl, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, CO<sub>2</sub>R<sup>3</sup>, CN, CF<sub>3</sub>, and halo.

3. The compound of claim 2 wherein W is selected from the group consisting of pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, optionally substituted with one or two substituents selected from the group consisting of optionally substituted C<sub>1-6</sub>alkyl, aryl, heteroaryl, N(R<sup>3</sup>)<sub>2</sub>, OR<sup>3</sup>, C(O)OR<sup>3</sup>, C(O)N(R<sup>3</sup>)<sub>2</sub>, and halo.


4. The compound of claim 2 wherein W is selected from the group consisting of




,




,




,




,



,




, and




- 79 -

5. The compound of claim 2 wherein W is selected from the group consisting of



and



6. The compound of claim 2 wherein W is pyrazinyl.







7. The compound of claim 1 wherein R<sup>6</sup> is OR<sup>11</sup>.

8. The compound of claim 1 wherein R<sup>7</sup> is heteroaryl.




- 80 -


9. The compound of claim 1 wherein the heteroaryl substituent on W and the heteroaryl group of R<sup>6</sup>, independently, are selected from the group consisting of




- 81 -




,




,




,




,




,




,



,












,








,


- 82 -








- 83 -




, and









11. A composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

12. A method of inhibiting checkpoint kinase 1 in a cell comprising a step of contacting the cell with an effective amount of a compound of claim 1.

13. A method of sensitizing cells in an individual undergoing a chemotherapeutic or radiotherapeutic treatment for a medical condition, comprising administering to the individual a therapeutically effective amount of a compound of claim 1 in combination with a chemotherapeutic agent, a radiotherapeutic agent, or a mixture thereof.

14. The method of claim 13 further comprising administering one or more cytokine, lymphokine, growth factor, or other hematopoietic factor.

15. The method of claim 12 wherein the chemotherapeutic agent is selected from the group consisting of an alkylating agent, an antimetabolite, a hormone or antagonist thereof, a radioisotope, an antibody, and mixtures thereof.

- 87 -

16. The method of claim 13 wherein the radio-therapeutic agent is selected from the group consisting of gamma-radiation, X-ray radiation, ultraviolet light, visible light, infrared radiation, and microwave radiation.

17. The method of claim 13 wherein the condition is a cancer selected from the group consisting of a colorectal cancer, a head and neck cancer, a pancreatic cancer, a breast cancer, a gastric cancer, a bladder cancer, a vulvar cancer, a leukemia, a lymphoma, a melanoma, a renal cell carcinoma, an ovarian cancer, a brain tumor, an osteosarcoma, and a lung carcinoma.



- 88 -

18. The method of claim 13 wherein the condition is a cancer selected from the group consisting of myxoid and round cell carcinomas, a locally advanced tumor, metastatic cancer, Ewing's sarcoma, a cancer metastase, a lymphatic metastase, squamous cell carcinoma, esophageal squamous cell carcinoma, oral carcinoma, multiple myeloma, acute lymphocytic leukemia, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer, small cell carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, nonsmall cell cancers, breast cancer, small cell carcinoma, ductal carcinoma, stomach cancer, colon cancer, colorectal cancer, polyps associated with colorectal neoplasia, pancreatic cancer, liver cancer, bladder cancer, primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, muscle-invasive bladder cancer, prostate cancer, ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid tumors in the ovarian follicle, testicular cancer, penile cancer, renal cell carcinoma, intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, osteomas and osteosarcomas, malignant melanoma, tumor progression of human skin keratinocytes, squamous cell cancer, thyroid cancer, retinoblastoma, neuroblastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma.

- 89 -

19. The method of claim 12 wherein the treatment is administered for an inflammatory condition selected from the group consisting of rheumatoid arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus erythematosus.

20. The method of claim 13 wherein the compound of claim 1 has at least a 20-fold selectivity in inhibiting Chk1 over protein kinase A, protein kinase C, cdc2, and pp60v-src.

21. The method of claim 13 wherein the compound of claim 1 has at least a 75-fold selectivity in inhibiting Chk1 over protein kinase A, protein kinase C, cdc2, and pp60v-src.

22. The method of claim 13 wherein the compound of claim 1 has at least a 100-fold selectivity in inhibiting Chk1 over protein kinase A, protein kinase C, cdc2, and pp60v-src.

23. The method of claim 13 wherein the chemotherapeutic agent comprises gemcitabine, pemetrexed, cisplatin, carboplatin, paclitaxel, or mixtures thereof.

24. A method of inhibiting aberrant cell proliferation comprising contacting a cell population comprising aberrantly proliferating cells with a Chk1 activator to substantially synchronize cell cycle arrest among said aberrantly proliferating cells, and subsequently contacting said cell population with a compound of claim 1 to substantially abrogate said cell cycle arrest.

- 90 -

25. The method of claim 24 wherein said Chk1 activator comprises at least one chemotherapeutic agent.

26. The method of claim 24 wherein said Chk1 activator comprises ionizing or ultraviolet radiation.

27. The method of claim 24 wherein said ionizing radiation is administered in conjunction with a radiosensitizer, a photosensitizer, or a mixture thereof.

28. The method of claim 24 wherein said aberrantly proliferating cells are noncancerous.

## SEQUENCE LISTING

&lt;110&gt; ICOS Corporation

&lt;120&gt; Compounds Useful for Inhibiting CHK1

&lt;130&gt; 27866/40321A

&lt;150&gt; US 60/602,968

&lt;151&gt; 2004-08-19

&lt;160&gt; 1

&lt;170&gt; PatentIn version 3.3

&lt;210&gt; 1

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; artificial

&lt;220&gt;

&lt;223&gt; synthetic peptide

&lt;400&gt; 1

Leu Tyr Arg Ser Pro Ser Met Pro Glu Asn Leu Asn Arg Arg Arg Arg  
1 5 10 15

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number  
**WO 2006/021002 A3**

(51) International Patent Classification:

C07D 241/20 (2006.01) A61K 31/4965 (2006.01)  
C07D 401/04 (2006.01) A61K 31/497 (2006.01)  
C07D 403/04 (2006.01) A61P 35/00 (2006.01)  
C07D 413/04 (2006.01)

(21) International Application Number:

PCT/US2005/029518

(22) International Filing Date: 18 August 2005 (18.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/602,968 19 August 2004 (19.08.2004) US

(71) Applicant (for all designated States except US): ICOS CORPORATION [US/US]; 22021 20th Avenue S.E., Bothell, WA 98021 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FAROUZ, Francine, S. [FR/US]; 4222 93rd Avenue S.E., Mercer Island, WA 98040 (US). HOLCOMB, Ryan [US/US]; 1214 2nd Avenue, Kirkland, WA 98033 (US). THORSETT, Eugene [US/US]; 410 2nd Ave S., #304, Kirkland, WA 98033 (US). GAUDINO, John, Joseph [US/US]; 4224 Prairie Fire Circle, Longmont, CO 80503 (US).

(74) Agent: NAPOLI, James, J.; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, Suite 6300, Sears Tower, Chicago, IL 60606-6357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
20 April 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS USEFUL FOR INHIBITING CHK1

(57) Abstract: Aryl- and heteroaryl-substituted urea compounds useful in the treatment of diseases and conditions related to DNA damage or lesions in DNA replication are disclosed. Methods of making the compounds, and their use as therapeutic agents, for example, in treating cancer and other diseases characterized by defects in DNA replication, chromosome segregation, or cell division also are disclosed.



## INTERNATIONAL SEARCH REPORT

/US2005/029518

## A. CLASSIFICATION OF SUBJECT MATTER

C07D241/20 C07D401/04 C07D403/04 C07D413/04 A61K31/4965  
 A61K31/497 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/070494 A (ICOS CORPORATION) 12 September 2002 (2002-09-12) cited in the application claims 1,2,6,8,9,14-18,28; compounds 313,232,299 page 31, line 19 - page 32, line 23 page 34, line 17 - line 21 page 47, line 18 - page 50, line 6	1-28
X	WO 03/093297 A (EXELIXIS, INC; BUHR, CHRIS, A; BAIK, TAE-GON) 13 November 2003 (2003-11-13)  claims 1,42,45-54; compound 919 paragraph '0032! - paragraph '0035! ----- -/--	1,7,9, 11-13, 15, 17-25,28

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

24 February 2006

Date of mailing of the international search report

09/03/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Krische, D

## INTERNATIONAL SEARCH REPORT

US2005/029518

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ARCADI A ET AL: "PALLADIUM-CATALYZED REACTION..." JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, US, vol. 61, 1996, pages 9280-9288, XP002073390 ISSN: 0022-3263 page 9288; compound 16A -----	1
X	NOMURA, MASAHIRO ET AL.: J. MED. CHEM., vol. 46, 2003, pages 3581-3599, XP002367530 table 2; compounds 38,40 -----	1,7,11
X	WO 99/32436 A (BAYER CORPORATION) 1 July 1999 (1999-07-01) abstract; claims 1,4 tables 2,4; compounds 47,48,51,52,55,131 -----	1,7,9,11
X	GB 921 682 A (J. R. GEIGY A.-G) 20 March 1963 (1963-03-20) claim 1 page 4, line 6 page 5, line 9 page 6, line 2 - line 6 page 6, line 9 - line 10 -----	1,7,11
X	DE 20 04 302 A1 (AGRIPAT S.A.) 6 August 1970 (1970-08-06) claims 1,10; table 1; compounds 8-11,21,25,26,29,37-39,57,61,66,81,84,85,8 9,98,100 -----	1,7
X	NAGAR, S. ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 16, 1973, pages 178-180, XP002367531 table 1; compounds 8,18 -----	1,7,11
X	US 4 071 524 A (BANITT ET AL) 31 January 1978 (1978-01-31) abstract; example 30; table 4 -----	1,7,11
X	VAN MUIJLWIJK-KOEZEN ET AL: "Isoquinoline and Quinazoline Urea Analogues as Antagonists for the Human Adenosine A3 Receptor" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 43, no. 5, 1 June 2000 (2000-06-01), pages 2227-2338, XP002147879 ISSN: 0022-2623 table 1; compounds 10A,11 -----	1,2,7,11

-/--



## INTERNATIONAL SEARCH REPORT

'US2005/029518

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 2005/027907 A (ICOS CORPORATION; CLARK, DARCEY; KEEGAN, KATHLEEN, S; PETERSON, SCOTT;) 31 March 2005 (2005-03-31) claims 1,2,11-66 page 138, line 6 - page 141, line 5 -----	1-28

## INTERNATIONAL SEARCH REPORT

PCT/US2005/029518

**Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 12-28 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-28 in part

Compounds having the general formula of claim 1 having  
X1=X2=N, Y=O or S, and W=CF<sub>3</sub>-substituted pyrazinyl for use  
as Chk1 inhibitors and compositions and methods using them.

---

2. claims: 1-28 in part

Compounds having the general formula of claim 1 having  
X1=X2=N, Y=O or S, and W=heteroaryl-substituted pyrazinyl  
for use as Chk1 inhibitors and compositions and methods  
using them.

---

## INTERNATIONAL SEARCH REPORT

Information on patent family members

'US2005/029518

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02070494	A	12-09-2002	BR 0207811 A	01-02-2005
			CA 2439709 A1	12-09-2002
			CN 1507436 A	23-06-2004
			EP 1379510 A1	14-01-2004
			HR 20030688 A2	31-10-2005
			JP 2004523568 T	05-08-2004
			MX PA03007920 A	04-12-2003
			NO 20033858 A	10-10-2003
			PL 364564 A1	13-12-2004
			ZA 200306721 A	03-05-2004
WO 03093297	A	13-11-2003	AU 2003234464 A1	17-11-2003
			CA 2484209 A1	13-11-2003
			EP 1501514 A2	02-02-2005
			JP 2005530760 T	13-10-2005
WO 9932436	A	01-07-1999	AT 291011 T	15-04-2005
			AU 763024 B2	10-07-2003
			AU 1905499 A	12-07-1999
			BG 64594 B1	31-08-2005
			BG 104599 A	30-03-2001
			BR 9814375 A	21-05-2002
			CA 2315646 A1	01-07-1999
			CN 1283180 A	07-02-2001
			DE 69829412 D1	21-04-2005
			DE 69829412 T2	28-07-2005
			DE 1049664 T1	03-05-2001
			EP 1049664 A1	08-11-2000
			ES 2153809 T1	16-03-2001
			GR 2001300006 T1	28-02-2001
			HU 0004437 A2	28-06-2001
			ID 26956 A	22-02-2001
			JP 2001526258 T	18-12-2001
			MX PA00006231 A	18-09-2002
			NO 20003230 A	21-08-2000
			NZ 505843 A	30-06-2003
			PL 342078 A1	21-05-2001
			SK 9612000 A3	12-03-2001
			TR 200002616 T2	21-11-2000
			TR 200100874 T2	21-06-2001
			UA 72451 C2	15-12-2000
GB 921682	A	20-03-1963	CH 362068 A	31-05-1962
			CH 364384 A	30-05-1962
			DE 1168606 B	23-04-1964
			ES 260391 A1	01-04-1961
			NL 254871 A	
			US 3230141 A	18-01-1966
DE 2004302	A1	06-08-1970	BE 745213 A1	30-07-1970
			CH 503460 A	13-11-1970
			CH 493499 A	15-07-1970
			CH 493480 A	15-07-1970
			CH 152169 D	13-11-1970
			ES 376044 A1	01-06-1972
			ES 376045 A1	16-04-1972
			ES 376046 A1	16-04-1972
			FR 2034515 A5	11-12-1970

# INTERNATIONAL SEARCH REPORT

Information on patent family members

'US2005/029518

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2004302	A1	GB 1270901 A	19-04-1972
		JP 48028897 B	05-09-1973
		NL 7001392 A	04-08-1970
US 4071524	A	31-01-1978	US 4154941 A
			15-05-1979
WO 2005027907	A	31-03-2005	NONE

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**